

References

- Abbott RA & Davies PS (2004) Habitual physical activity and physical activity intensity: their relation to body composition in 5.0-10.5-y-old children. *Eur J Clin Nutr* **58**, 285-291.
- Aleman-Mateo H, Salazar G, Hernandez-Triana M & Valencia ME (2006) Total energy expenditure, resting metabolic rate and physical activity level in free-living rural elderly men and women from Cuba, Chile and Mexico. *Eur J Clin Nutr*.
- Amatruda JM, Statt MC & Welle SL (1993) Total and resting energy expenditure in obese women reduced to ideal body weight. *J Clin Invest* **92**, 1236-1242.
- Anderson SE, Bandini LG, Dietz WH & Must A (2004) Relationship between temperament, nonresting energy expenditure, body composition, and physical activity in girls. *Int J Obes Relat Metab Disord* **28**, 300-306.
- Andersen LF, Tomten H, Haggarty P, Lovo A & Hustvedt BE (2003) Validation of energy intake estimated from a food frequency questionnaire: a doubly labelled water study. *Eur J Clin Nutr* **57**, 279-284.
- Arvidsson D, Slinde F & Hulthen L (2005) Physical activity questionnaire for adolescents validated against doubly labelled water. *Eur J Clin Nutr* **59**, 376-383.
- Ball EJ, O'Connor J, Abbott R, Steinbeck KS, Davies PS, Wishart C, Gaskin KJ & Baur LA (2001) Total energy expenditure, body fatness, and physical activity in children aged 6-9 y. *Am J Clin Nutr* **74**, 524-528.
- Bandini LG, Must A, Spadano JL & Dietz WH (2002) Relation of body composition, parental overweight, pubertal stage, and race-ethnicity to energy expenditure among premenarcheal girls. *Am J Clin Nutr* **76**, 1040-1047.
- Bandini LG, Schoeller DA & Dietz WH (1990b) Energy expenditure in obese and nonobese adolescents. *Pediatr Res* **27**, 198-203
- Black AE, Bingham SA, Johansson G & Coward WA (1997) Validation of dietary intakes of protein and energy against 24 hour urinary N and DLW energy expenditure in middle-aged women, retired men and post-obese subjects: comparisons with validation against presumed energy requirements. *Eur J Clin Nutr* **51**, 405-413.
- Black AE, Welch AA & Bingham SA (2000) Validation of dietary intakes measured by diet history against 24h urinary nitrogen excretion and energy expenditure measured by DLW method in middle-aged women. *Br J Nutr* **83**, 341-354.
- Blanc S, Schoeller DA, Bauer D, Danielson ME, Tylavsky F, Simonsick EM, Harris TB, Kritchevsky SB & Everhart JE (2004) Energy requirements in the eighth decade of life. *Am J Clin Nutr* **79**, 303-310.
- Blanton CA, Moshfegh AJ, Baer DJ & Kretsch MJ (2006) The USDA Automated

- Multiple-Pass Method Accurately Estimates Group Total Energy and Nutrient Intake. *J Nutr* **136**, 2594-2599.
- Bratteby LE, Sandhagen B, Fan H & Samuelson G (1997) A 7-day activity diary for assessment of daily energy expenditure validated by the doubly labelled water method in adolescents. *Eur J Clin Nutr* **51**, 585-591.
- Bratteby LE, Sandhagen B & Samuelson G (2005) Physical activity, energy expenditure and their correlates in two cohorts of Swedish subjects between adolescence and early adulthood. *Eur J Clin Nutr* **59**, 1324-1334.
- Brochu M, Starling RD, Ades PA & Poehlman ET (1999) Are aerobically fit older individuals more physically active in their free-living time? A doubly labeled water approach. *J Clin Endocrinol Metab* **84**, 3872-3876.
- Butte NF, Wong WW & Hopkinson JM (2001) Energy requirements of lactating women derived from doubly labeled water and milk energy output. *J Nutr* **131**, 53-58.
- Butte NF, Wong WW, Hopkinson JM, Heinz CJ, Mehta NR & Smith EO (2000) Energy requirements derived from total energy expenditure and energy deposition during the first 2 y of life. *Am J Clin Nutr* **72**, 1558-1569.
- Butte NF, Wong WW, Treuth MS, Ellis KJ & O'Brian Smith E (2004) Energy requirements during pregnancy based on total energy expenditure and energy deposition. *Am J Clin Nutr* **79**, 1078-1087.
- Carpenter WH, Fonong T, Toth MJ, Ades PA, Calles-Escandon J, Walston JD & Poehlman ET (1998) Total daily energy expenditure in free-living older African-Americans and Caucasians. *Am J Physiol* **274**, E96-101.
- Champagne CM, Baker NB, Delany JP, Harsha DW & Bray GA (1998) Assessment of energy intake underreporting by doubly labeled water and observations on reported nutrient intakes in children. *J Am Diet Assoc* **98**, 426-433.
- Champagne CM, Bray GA, Kurtz AA, Monteiro JB, Tucker E, Volaufova J & DeLany JP (2002) Energy intake and energy expenditure: a controlled study comparing dietitians and non-dietitians. *J Am Diet Assoc* **102**, 1428-1432.
- Chong PK, Jung RT, Rennie MJ & Scrimgeour CM (1993) Energy expenditure in lean and obese diabetic patients using the doubly labelled water method. *Diabet Med* **10**, 729-735.
- Clark D, Tomas F, Withers RT, Chandler C, Brinkman M, Phillips J, Berry M, Ballard FJ & Nestel P (1994) Energy metabolism in free-living, 'large-eating' and 'small-eating' women: studies using 2H₂(18)O. *Br J Nutr* **72**, 21-31.
- Conway JM, Seale JL, Jacobs DR, Jr., Irwin ML & Ainsworth BE (2002) Comparison of energy expenditure estimates from doubly labeled water, a physical activity questionnaire, and physical activity records. *Am J Clin Nutr* **75**, 519-525.
- Coward WA, Wright A & Bluck LJC. Comparison between energy intake and energy

expenditure in the NDNS survey, adults aged 19-64 years. MRC Resource Centre for Human Nutrition Research (Cambridge, 2000). Paper produced from the feasibility study data for NDNS steering committee.

- Craig SB, Bandini LG, Lichtenstein AH, Schaefer EJ & Dietz WH (1996) The impact of physical activity on lipids, lipoproteins, and blood pressure in preadolescent girls. *Pediatrics* **98**, 389-395.
- Davidson L, McNeill G, Haggarty P, Smith JS & Franklin MF (1997) Free-living energy expenditure of adult men assessed by continuous heart-rate monitoring and doubly-labelled water. *Br J Nutr* **78**, 695-708.
- Davies PS, Ewing G & Lucas A (1989) Energy expenditure in early infancy. *Br J Nutr* **62**, 621-629.
- Davies PS, Gregory J & White A (1995) Energy expenditure in children aged 1.5 to 4.5 years: a comparison with current recommendations for energy intake. *Eur J Clin Nutr* **49**, 360-364.
- Davies PS, Wells JC, Hinds A, Day JM & Laidlaw A (1997) Total energy expenditure in 9 month and 12 month infants. *Eur J Clin Nutr* **51**, 249-252.
- DeLany JP, Bray GA, Harsha DW & Volaufova J (2002) Energy expenditure in preadolescent African American and white boys and girls: the Baton Rouge Children's Study. *Am J Clin Nutr* **75**, 705-713.
- DeLany JP, Bray GA, Harsha DW & Volaufova J (2004) Energy expenditure in African American and white boys and girls in a 2-y follow-up of the Baton Rouge Children's Study. *Am J Clin Nutr* **79**, 268-273.
- DeLany JP, Bray GA, Harsha DW & Volaufova J (2006) Energy expenditure and substrate oxidation predict changes in body fat in children. *Am J Clin Nutr* **84**, 862-870.
- Ekelund U, Aman J, Yngve A, Renman C, Westerterp K & Sjostrom M (2002) Physical activity but not energy expenditure is reduced in obese adolescents: a case-control study. *Am J Clin Nutr* **76**, 935-941.
- Ekelund U, Sjostrom M, Yngve A, Poortvliet E, Nilsson A, Froberg K, Wedderkopp N & Westerterp K (2001) Physical activity assessed by activity monitor and doubly labeled water in children. *Med Sci Sports Exerc* **33**, 275-281.
- Ekelund U, Yngve A, Brage S, Westerterp K & Sjostrom M (2004) Body movement and physical activity energy expenditure in children and adolescents: how to adjust for differences in body size and age. *Am J Clin Nutr* **79**, 851-856.
- Fogelholm M, Hiilloskorpi H, Laukkanen R, Oja P, Van Marken LW & Westerterp K (1998) Assessment of energy expenditure in overweight women. *Med Sci Sports Exerc* **30**, 1191-1197
- Fontvieille AM, Harper IT, Ferraro RT, Spraul M & Ravussin E (1993) Daily energy expenditure by five-year-old children, measured by doubly labelled water. *J*

Pediatr **123**, 200-207.

- Forsum E, Kabir N, Sadurskis A & Westerterp K (1992) Total energy expenditure of healthy Swedish women during pregnancy and lactation. *Am J Clin Nutr* **56**, 334-342.
- Franks PW, Ravussin E, Hanson RL, Harper IT, Allison DB, Knowler WC, Tataranni PA & Salbe AD (2005) Habitual physical activity in children: the role of genes and the environment. *Am J Clin Nutr* **82**, 901-908.
- Fuller NJ, Sawyer MB, Coward WA, Paxton P & Elia M (1996) Components of total energy expenditure in free-living elderly men (over 75 years of age): measurement, predictability and relationship to quality-of-life indices. *Br J Nutr* **75**, 161-173.
- Gibney ER, Murgatroyd P, Wright A, Jebb S & Elia M (2003) Measurement of total energy expenditure in grossly obese women: comparison of the bicarbonate-urea method with whole-body calorimetry and free-living doubly labelled water. *Int J Obes Relat Metab Disord* **27**, 641-647.
- Goldberg GR, Prentice AM, Coward WA, Davies HL, Murgatroyd PR, Sawyer MB, Ashford J & Black AE (1991) Longitudinal assessment of the components of energy balance in well-nourished lactating women. *Am J Clin Nutr* **54**, 788-798.
- Goldberg GR, Prentice AM, Coward WA, Davies HL, Murgatroyd PR, Wensing C, Black AE, Harding M & Sawyer M (1993) Longitudinal assessment of energy expenditure in pregnancy by the doubly labeled water method. *Am J Clin Nutr* **57**, 494-505.
- Goran MI, Calles-Escandon J, Poehlman ET, O'Connell M & Danforth E Jr (1994) Effects of increased energy intake and/or physical activity on energy expenditure in young healthy men. *J Appl Physiol* **77**, 366-372.
- Goran MI, Carpenter WH, McGloin A, Johnson R, Hardin JM & Weinsier RL (1995a) Energy expenditure in children of lean and obese parents. *Am J Physiol* **268**, E917-E924.
- Goran MI, Carpenter WH & Poehlman ET (1993b) Total energy expenditure in 4- to 6-yr-old children. *Am J Physiol* **264**, E706-E711.
- Goran MI, Gower BA, Nagy TR & Johnson RK (1998a) Developmental changes in energy expenditure and physical activity in children: evidence for a decline in physical activity in girls before puberty. *Pediatrics* **101**, 887-891.
- Goran MI & Poehlman ET (1992) Total energy expenditure and energy requirements in healthy elderly persons. *Metabolism* **41**, 744-753.
- Haggarty P, McNeill G, Manneh MK, Davidson L, Milne E, Duncan G & Ashton J (1994) The influence of exercise on the energy requirements of adult males in the UK. *Br J Nutr* **72**, 799-813.

- Hernandez-Triana M, Salazar G, Diaz E, Sanchez V, Basabe B, Gonzalez S & Diaz ME (2002) Total energy expenditure by the doubly-labeled water method in rural preschool children in Cuba. *Food Nutr Bull* **23**, 76-81.
- Hibbert JM, Broemeling LD, Isenberg JN & Wolfe RR (1994) Determinants of free-living energy expenditure in normal weight and obese women measured by doubly labeled water. *Obes Res* **2**, 44-53
- Hoffman DJ, Sawaya AL, Coward WA, Wright A, Martins PA, de Nascimento C, Tucker KL & Roberts SB (2000) Energy expenditure of stunted and nonstunted boys and girls living in the shantytowns of Sao Paulo, Brazil. *Am J Clin Nutr* **72**, 1025-1031.
- Hoos MB, Plasqui G, Gerver WJ & Westerterp KR (2003b) Physical activity level measured by doubly labeled water and accelerometry in children. *Eur J Appl Physiol* **89**, 624-626.
- Hunter GR, Weinsier RL, Darnell BE, Zuckerman PA & Goran MI (2000a) Racial differences in energy expenditure and aerobic fitness in premenopausal women. *Am J Clin Nutr* **71**, 500-506.
- Irwin ML, Ainsworth BE & Conway JM (2001) Estimation of energy expenditure from physical activity measures: determinants of accuracy. *Obes Res* **9**, 517-525.
- Jiang Z, Yan Q, Su Y, Acheson KJ, Thelin A, Pigué-Welsch C, Ritz P & Ho ZC (1998) Energy expenditure of Chinese infants in Guangdong Province, south China, determined with use of the doubly labeled water method. *Am J Clin Nutr* **67**, 1256-1264.
- Johnson RK, Russ J & Goran MI (1998) Physical activity related energy expenditure in children by doubly labeled water as compared with the Caltrac accelerometer. *Int J Obes Relat Metab Disord* **22**, 1046-1052.
- Jones PJ, Martin LJ, Su W & Boyd NF (1997) Canadian recommended nutrient intakes underestimate true energy requirements in middle-aged women. *Can J Public Health* **88**, 314-319.
- Kaczkowski CH, Jones PJ, Feng JY & Bayley HS (2000b) Canadian recommendations underestimate energy needs of women over fifty years as determined by doubly-labelled water. *Can J Physiol Pharmacol* **78**, 631-635.
- Kroke A, Klipstein-Grobusch K, Voss S, Mosender J, Thielecke F, Noack R & Boeing H (1999) Validation of a self administered food frequency questionnaire administered in the European Prospective Investigation into Cancer and Nutrition (EPIC) study: comparison of energy, protein and macronutrient intakes estimated with doubly labelled water, urinary nitrogen and repeated 24-h dietary recall methods. *Am J Clin Nutr* **70**: 439-47.
- Kushner RF, Racette SB, Neil K & Schoeller DA (1995) Measurement of physical activity among black and white obese women. *Obes Res* **3 Suppl 2**, 261s-

265s.

- Lanigan JA, Wells JC, Lawson MS & Lucas A (2001) Validation of food diary method for assessment of dietary energy and macronutrient intake in infants and children aged 6-24 months. *Eur J Clin Nutr* **55**, 124-129.
- Lucas A, Ewing G, Roberts SB & Coward WA (1987) How much energy does the breast fed infant consume and expend? *Br Med J (Clin Res Ed)* **295**, 75-77.
- Leenders NYJM, Sherman WM, Nagaraja HN & Kien CL (2001) Evaluation of methods to assess physical activity in free living conditions. *Med Sci Sports Exerc* **33**, 1233-1240.
- Lindquist CH, Cummings T & Goran MI (2000) Use of tape-recorded food records in assessing children's dietary intake. *Obes Res* **8**, 2-11.
- Livingstone MB, Coward WA, Prentice AM, Davies PS, Strain JJ, McKenna PG, Mahoney CA, White JA, Stewart CM & Kerr MJ (1992a) Daily energy expenditure in free-living children: comparison of heart-rate monitoring with the doubly labeled water (2H₂(18)O) method. *Am J Clin Nutr* **56**, 343-352.
- Livingstone MB, Prentice AM, Coward WA, Ceesay SM, Strain JJ, McKenna PG, Nevin GB, Barker ME & Hickey RJ (1990) Simultaneous measurement of free-living energy expenditure by the doubly labeled water method and heart-rate monitoring. *Am J Clin Nutr* **52**, 59-65.
- Lof M & Forsum E (2004) Validation of energy intake by dietary recall against different methods to assess energy expenditure. *J Hum Nutr Diet* **17**, 471-480.
- Lof M & Forsum E (2006) Activity pattern and energy expenditure due to physical activity before and during pregnancy in healthy Swedish women. *Br J Nutr* **95**, 296-302.
- Lof M, Hannestad U & Forsum E (2003) Comparison of commonly used procedures, including the doubly-labelled water technique, in the estimation of total energy expenditure of women with special reference to the significance of body fatness. *Br J Nutr* **90**, 961-968.
- Lopez-Alarcon M, Merrifield J, Fields DA, Hilario-Hailey T, Franklin FA, Shewchuk RM, Oster RA & Gower BA (2004) Ability of the actiwatch accelerometer to predict free-living energy expenditure in young children. *Obes Res* **12**, 1859-1865.
- Lovelady CA, Meredith CN, McCrory MA, Nommsen LA, Joseph LJ & Dewey KG (1993) Energy expenditure in lactating women: a comparison of doubly labeled water and heart-rate-monitoring methods. *Am J Clin Nutr* **57**, 512-518.
- Maffeis C, Pinelli L, Zaffanello M, Schena F, Iacumin P & Schutz Y (1995) Daily energy expenditure in free-living conditions in obese and non-obese children: comparison of doubly labelled water (2H₂(18)O) method and heart-rate monitoring. *Int J Obes Relat Metab Disord* **19**, 671-677.

- Manini TM, Everhart JE, Patel KV, Schoeller DA, Colbert LH, Visser M, Tylavsky F, Bauer DC, Goodpaster BH & Harris TB (2006) Daily Activity Energy Expenditure and Mortality Among Older Adults. *JAMA* **296**, 171-179.
- Masse LC, Fulton JE, Watson KL, Mahar MT, Meyers MC & Wong WW (2004) Influence of body composition on physical activity validation studies using doubly labeled water. *J Appl Physiol* **96**, 1357-1364.
- McGloin AF, Livingstone MB, Greene LC, Webb SE, Gibson JM, Jebb SA, Cole TJ, Coward WA, Wright A & Prentice AM (2002) Energy and fat intake in obese and lean children at varying risk of obesity. *Int J Obes Relat Metab Disord* **26**, 200-207.
- Meijer GA, Westerterp KR, van Hulsel AM & Ten Hoor F (1992) Physical activity and energy expenditure in lean and obese adult human subjects. *Eur J Appl Physiol Occup Physiol* **65**, 525-528.
- Melby CL, Ho RC, Jeckel K, Beal L, Goran M & Donahoo WT (2000) Comparison of risk factors for obesity in young, nonobese African-American and Caucasian women. *Int J Obes Relat Metab Disord* **24**, 1514-1522.
- Montgomery C, Reilly JJ, Jackson DM, Kelly LA, Slater C, Paton JY & Grant S (2004) Relation between physical activity and energy expenditure in a representative sample of young children. *Am J Clin Nutr* **80**, 591-596.
- Montgomery C, Reilly JJ, Jackson DM, Kelly LA, Slater C, Paton JY & Grant S (2005) Validation of energy intake by 24-hour multiple pass recall: comparison with total energy expenditure in children aged 5-7 years. *Br J Nutr* **93**, 671-676.
- Morio B, Ritz P, Verdier E, Montaurier C, Beaufriere B & Vermorel M (1997) Critical evaluation of the factorial and heart-rate recording methods for the determination of energy expenditure of free-living elderly people. *Br J Nutr* **78**, 709-722.
- Nagy TR, Gower BA, Shewchuk RM & Goran MI (1997) Serum leptin and energy expenditure in children. *J Clin Endocrinol Metab* **82**, 4149-4153.
- Nguyen VT, Larson DE, Johnson RK & Goran MI (1996) Fat intake and adiposity in children of lean and obese parents. *Am J Clin Nutr* **63**, 507-513.
- O'Connor J, Ball EJ, Steinbeck KS, Davies PS, Wishart C, Gaskin KJ & Baur LA (2001) Comparison of total energy expenditure and energy intake in children aged 6-9 y. *Am J Clin Nutr* **74**, 643-649.
- Pannemans DL & Westerterp KR (1995) Energy expenditure, physical activity and basal metabolic rate of elderly subjects. *Br J Nutr* **73**, 571-581.
- Papamandjaris AA, White MD & Jones PJ (1999) Components of total energy expenditure in healthy young women are not affected after 14 days of feeding with medium-versus long-chain triglycerides. *Obes Res* **7**, 273-280.

- Paul DR, Novotny JA & Rumpler WV (2004) Effects of the interaction of sex and food intake on the relation between energy expenditure and body composition. *Am J Clin Nutr* **79**, 385-389.
- Perks SM, Roemmich JN, Sadow-Pajewski M, Clark PA, Thomas E, Weltman A, Patrie J & Rogol AD (2000) Alterations in growth and body composition during puberty. IV. Energy intake estimated by the Youth-Adolescent Food-Frequency Questionnaire: validation by the doubly labeled water method. *Am J Clin Nutr* **72**, 1455-1460.
- Philippaerts RM, Westerterp KR & Lefevre J (1999) Doubly labelled water validation of three physical activity questionnaires. *Int J Sports Med* **20**, 284-289
- Platte P, Pirke KM, Trimborn P, Pietsch K, Krieg JC & Fichter MM (1994) Resting metabolic rate and total energy expenditure in acute and weight recovered patients with anorexia nervosa and in healthy young women. *Int J Eat Disord* **16**, 45-52.
- Prentice AM, Black AE, Coward WA, Davies HL, Goldberg GR, Murgatroyd PR, Ashford J, Sawyer M & Whitehead RG (1986) High levels of energy expenditure in obese women. *Br Med J (Clin Res Ed)* **292**, 983-987.
- Reilly JJ, Lord A, Bunker VW, Prentice AM, Coward WA, Thomas AJ & Briggs RS (1993) Energy balance in healthy elderly women. *Br J Nutr* **69**, 21-27.
- Reilly JJ, Montgomery C, Jackson D, MacRitchie J & Armstrong J (2001) Energy intake by multiple pass 24 h recall and total energy expenditure: a comparison in a representative sample of 3-4-year-olds. *Br J Nutr* **86**, 601-605.
- Rennie KL, Livingstone MB, Wells JC, McGloin A, Coward WA, Prentice AM & Jebb SA (2005) Association of physical activity with body-composition indexes in children aged 6-8 y at varied risk of obesity. *Am J Clin Nutr* **82**, 13-20.
- Roberts SB, Heyman MB, Evans WJ, Fuss P, Tsay R & Young VR (1991) Dietary energy requirements of young adult men, determined by using the doubly labeled water method. *Am J Clin Nutr* **54**, 499-505.
- Roberts SB, Young VR, Fuss P, Heyman MB, Fiatarone M, Dallal GE, Cortiella J & Evans WJ (1992) What are the dietary energy needs of elderly adults? *Int J Obes Relat Metab Disord* **16**, 969-976.
- Roemmich JN, Clark PA, Walter K, Patrie J, Weltman A & Rogol AD (2000) Pubertal alterations in growth and body composition. V. Energy expenditure, adiposity, and fat distribution. *Am J Physiol Endocrinol Metab* **279**, E1426-E1436.
- Rosetta L, Kurpad A, Mascie-Taylor CG & Shetty PS (2005) Total energy expenditure (H218O), physical activity level and milk output of lactating rural Bangladeshi tea workers and nontea workers. *Eur J Clin Nutr* **59**, 632-638.
- Rothenberg EM (2002) Resting, activity and total energy expenditure at age 91-96

compared to age 73. *J Nutr Health Aging* **6**, 177-178.

Rothenberg EM, Bosaeus IG, Westertep KR & Steen BC (2000) Resting energy expenditure, activity energy expenditure and total energy expenditure at age 91-96 years. *Br J Nutr* **84**, 319-324.

Rush EC, Plank LD & Coward WA (1999) Energy expenditure of young Polynesian and European women in New Zealand and relations to body composition. *Am J Clin Nutr* **69**, 43-48.

Rush EC, Plank LD, Davies PS, Watson P & Wall CR (2003) Body composition and physical activity in New Zealand Maori, Pacific and European children aged 5-14 years. *Br J Nutr* **90**, 1133-1139.

Salazar G, Vio F, Garcia C, Aguirre E & Coward WA (2000) Energy requirements in Chilean infants. *Arch Dis Child Fetal Neonatal Ed* **83**, F120-F123.

Seale JL (2002) Predicting total energy expenditure from self-reported dietary records and physical characteristics in adult and elderly men and women. *Am J Clin Nutr* **76**, 529-534.

Spadano JL, Bandini LG, Must A, Dallal GE & Dietz WH (2005) Longitudinal changes in energy expenditure in girls from late childhood through midadolescence. *Am J Clin Nutr* **81**, 1102-1109.

Starling RD, Matthews DE, Ades PA & Poehlman ET (1999) Assessment of physical activity in older individuals: a doubly labeled water study. *J Appl Physiol* **86**, 2090-2096.

Starling RD, Toth MJ, Matthews DE & Poehlman ET (1998) Energy requirements and physical activity of older free living African-Americans: a doubly labeled water study. *J Clin Endocrinol Metab* **83**, 1529-1534.

Stunkard AJ, Berkowitz RI, Stallings VA & Schoeller DA (1999) Energy intake, not energy output, is a determinant of body size in infants. *Am J Clin Nutr* **69**, 524-530.

Sun M, Gower BA, Nagy TR, Bartolucci AA & Goran MI (1999) Do hormonal indices of maturation explain energy expenditure differences in African American and Caucasian prepubertal children? *Int J Obes Relat Metab Disord* **23**, 1320-1326.

Sun M, Gower BA, Nagy TR, Trowbridge CA, Dezenberg C & Goran MI (1998) Total, resting, and activity-related energy expenditures are similar in Caucasian and African-American children. *Am J Physiol* **274**, E232-E237.

Tennefors C, Coward WA, Hernell O, Wright A & Forsum E (2003) Total energy expenditure and physical activity level in healthy young Swedish children 9 or 14 months of age. *Eur J Clin Nutr* **57**, 647-653.

Tomoyasu NJ, Toth MJ & Poehlman ET (1999) Misreporting of total energy intake in older men and women. *J Am Geriatr Soc* **47**, 710-715.

- Tomoyasu NJ, Toth MJ & Poehlman ET (2000) Misreporting of total energy intake in older African Americans. *Int J Obes Relat Metab Disord* **24**, 20-26.
- Tran KM, Johnson RK, Soultanakis RP & Matthews DE (2000) In-person vs telephone-administered multiple-pass 24-hour recalls in women: validation with doubly labeled water. *J Am Diet Assoc* **100**, 777-783.
- Treuth MS, Butte NF & Wong WW (2000) Effects of familial predisposition to obesity on energy expenditure in multiethnic prepubertal girls. *Am J Clin Nutr* **71**, 893-900.
- Treuth MS, Figueroa-Colon R, Hunter GR, Weinsier GR, Butte NF & Goran MI (1998) Energy expenditure and physical fitness in overweight vs non-overweight prepubertal girls. *Int J Obes* **22**, 440-447.
- Trowbridge CA, Gower BA, Nagy TR, Hunter GR, Treuth MS & Goran MI (1997) Maximal aerobic capacity in African-American and Caucasian prepubertal children. *Am J Physiol* **273**, E809-E814.
- Vasquez F, Salazar G, Andrade M, Vasquez L & Diaz E (2006) Energy balance and physical activity in obese children attending day-care centres. *Eur J Clin Nutr* **60**, 1115-1121.
- Velthuis-te Wierik EJ, Westerterp KR & van den BH (1995) Impact of a moderately energy-restricted diet on energy metabolism and body composition in non-obese men. *Int J Obes Relat Metab Disord* **19**, 318-324.
- Vinken AG, Bathalon GP, Sawaya AL, Dallal GE, Tucker KL & Roberts SB (1999) Equations for predicting the energy requirements of healthy adults aged 18-81 y. *Am J Clin Nutr* **69**, 920-926.
- Votruba SB, Blanc S & Schoeller DA (2002) Pattern and cost of weight gain in previously obese women. *Am J Physiol Endocrinol Metab* **282**, E923-E930.
- Walsh MC, Hunter GR, Sirikul B & Gower BA (2004) Comparison of self-reported with objectively assessed energy expenditure in black and white women before and after weight loss. *Am J Clin Nutr* **79**, 1013-1019.
- Warwick PM (2006) Factorial estimation of daily energy expenditure using a simplified method was improved by adjustment for excess post-exercise oxygen consumption and thermic effect of food. *Eur J Clin Nutr* **60**, 1337-1340.
- Washburn RA, Jacobsen DJ, Sonko BJ, Hill JO & Donnelly JE (2003) The validity of the Stanford Seven-Day Physical Activity Recall in young adults. *Med Sci Sports Exerc* **35**, 1374-1380
- Weber JL, Reid PM, Greaves KA, DeLany JP, Stanford VA, Going SB, Howell WH & Houtkooper LB (2001) Validity of self-reported energy intake in lean and obese young women, using two nutrient databases, compared with total energy expenditure assessed by doubly labeled water. *Eur J Clin Nutr* **55**, 940-950.

- Weinsier RL, Hunter GR, Zuckerman PA, Redden DT, Darnell BE, Larson DE, Newcomer BR & Goran MI (2000) Energy expenditure and free-living physical activity in black and white women: comparison before and after weight loss. *Am J Clin Nutr* **71**, 1138-1146.
- Wells JC, Cole TJ & Davies PS (1996) Total energy expenditure and body composition in early infancy. *Arch Dis Child* **75**, 423-426.
- Wells JC & Davies PSW (1995) Energy cost of physical activity in twelve week old infants. *Am J Hum Biol* **7**, 85-92.
- Welle S, Forbes GB, Statt M, Barnard RR & Amatruda JM (1992) Energy expenditure under free-living conditions in normal-weight and overweight women. *Am J Clin Nutr* **55**, 14-21.
- Westerterp KR, Meijer GA, Saris WH, Soeters PB, Winants Y & Ten Hoor F (1991) Physical activity and sleeping metabolic rate. *Med Sci Sports Exerc* **23**, 166-170.
- Withers RT, Smith DA, Tucker RC, Brinkman M & Clark DG (1998) Energy metabolism in sedentary and active 49- to 70-yr-old women. *J Appl Physiol* **84**, 1333-1340.
- Wong WW (1994) Energy expenditure of female adolescents. *J Am Coll Nutr* **13**, 332-337.
- Wong WW, Butte NF, Ellis KJ, Hergenroeder AC, Hill RB, Stuff JE & Smith EO (1999) Pubertal African-American girls expend less energy at rest and during physical activity than Caucasian girls. *J Clin Endocrinol Metab* **84**, 906-911.
- Yao M, McCrory MA, Ma G, Li Y, Dolnikowski GG & Roberts SB (2002) Energy requirements of urban Chinese adults with manual or sedentary occupations, determined using the doubly labeled water method. *Eur J Clin Nutr* **56**, 575-584.