

STATEMENT ON DIETARY FIBRE

August 2008

1. Draft SACN statement on dietary fibre and health and the dietary fibre definition

1.1 Current definitions

1. The current UK definition of dietary fibre (Department of Health 1991) is that dietary fibre should be defined as non starch polysaccharides (NSP) where this refers to non alpha-glucans as measured by the technique of Englyst and Cummings or other comparable techniques.
2. In 1997 the Committee for the Medical Aspects of Food Policy (COMA) subgroup on Dietary Fibre for Labelling purposes were unable to reach a consensus on a definition of dietary fibre for nutrition labelling purposes.

1.2 SACN statement on the definition for dietary fibre

3. SACN have reviewed the available scientific evidence for components under consideration for inclusion in the Codex definition on dietary fibre for nutrition labelling purposes, using the SACN frame work for assessing evidence (SACN 2002). These components include total fibre, non starch polysaccharides, fibre components from cereal, fibre components from fruit and vegetables, starch, resistant starch, polydextrose, oligosaccharides (including fructooligosaccharides (FOS), galactooligosaccharides (GOS) and inulin), lignin, soluble fibres (including pectin and guar gum).
4. There is some evidence to show that 'dietary fibre' (in the form of NSP) as an integral part of a balanced diet, high in food sources such as fruit and vegetables and cereals, appears to have positive effects on health. There are two potential explanations for this observation; firstly dietary fibre could be a surrogate marker for another component of the diet or, secondly, it could be that dietary fibre itself is having a direct effect.
5. Dietary components extracted from foods or synthesised chemically need to show clear evidence of a physiological effect before they can be assumed to be as effective as whole foods intrinsically containing fibres. It still needs to be proven that specific fibres components such as oligosaccharides have an independent effect, The working group

therefore consider that it is potentially misleading to include non intrinsic fibre in the definition in the absence of other evidence.

6. SACN consider that a material can be considered as dietary fibre if it is resistant to digestion and absorption in the small intestine and has a demonstrable physiological effect potentially associated with health benefits in the body, such as increasing stool bulk, decreasing intestinal transit time or decreasing post prandial glycaemia. Evidence only of increased fermentation in the gut should not be included under this definition, since although this has a direct effect on the microflora, it must also be shown to have a demonstrable benefit to the host to be considered as dietary fibre.
7. Based on the available evidence SACN conclude that there is sufficient evidence for an association between those compounds identified as NSP and colonic function (including stool weight/mass and transit time) and between those compounds identified as soluble fibre (from oats, psyllium, pectin and guar gum) and lowering of total cholesterol and LDL cholesterol. The committee note that there is insufficient evidence for health effects in relation to dietary fibre and colorectal cancer or colorectal adenoma, obesity, diabetes, blood pressure lowering and for prebiotic effects.
8. Taking the definition recommended by SACN these two fractions (NSP and soluble fibre) would be the only components definitely included within a fibre definition without further evidence. For other components to be considered for inclusion in the definition the proposed physiological effects need to be demonstrated.

2. Summary of evidence considered by SACN on fibre and health

9. A summary of the evidence considered by SACN and the conclusions drawn are presented below. A full review of this evidence will be included in a report on carbohydrate and health, which will be published when the committee have finished their deliberations.

2.1 Obesity and metabolic health

2.1.1 Observational studies

10. On balance, evidence from observational studies is suggestive that increasing amounts of total fibre (as determined by AOAC method) in the diet are associated with lower body weight and waist circumference. The single study (Iqbal *et al* 2006) to find no association investigated a smaller sample than the majority of studies where a positive association was seen. There is no evidence for an association of fibre intake on weight change in children.
11. Evidence from 12 prospective studies investigating total fibre intake and type 2 diabetes incidences is variable (Colditz *et al* 1992; Hodge *et al* 2004; Hu *et al* 2001; Krishnan *et al* 2007; Lindstrom *et al* 2006a; Lindstrom *et al* 2006b; Meyer *et al* 2000; Montonen *et al* 2003; Salmeron *et al* 1997a; Salmeron *et al* 1997b; Schulze *et al* 2004; Schulze *et al* 2007; Stevens *et al* 2002). The majority of studies have not seen any significant association of fibre with risk of diabetes. A number of the studies have investigated relationships with different sources of fibre. There is evidence of an association with

cereal fibre intake, with nine of eleven studies observing a significant reduction in diabetes incidence with increasing intake.

12. The totality of the evidence does not support an association for fibre from fruit, vegetables or legumes on the outcomes investigated. Two of three studies observed an association with insoluble fibre intakes, but no association with soluble fibre (Meyer *et al* 2000; Montonen *et al* 2003).
13. Amongst cross-sectional studies investigating the association of metabolic risk factors with fibre intake, a decreased risk of metabolic syndrome has been observed with increased cereal fibre intake, but not with total, fruit, vegetable or legume fibre (Mckeown *et al* 2004; Newby *et al* 2007). This is consistent with the evidence discussed for diabetes risk. Studies have investigated the relationship between fibre intake and a variety of measures of insulin sensitivity and glucose tolerance. Although not wholly consistent, these are suggestive of an association, particularly for total and cereal fibre.

2.1.2 Intervention Studies

14. Intervention studies have investigated the relationship between non-starch polysaccharide and food sources of fibre on weight and insulin sensitivity/ glucose tolerance ¹. The balance of evidence suggests that fibre supplementation with certain fibres in sufficient amounts is more likely to be efficacious in assisting weight loss as an adjunct to a weight-reducing diet than an *ad libitum* habitual diet. There is no evidence to suggest an association of fibre supplementation on weight control in children, which is consistent with evidence from observational studies.
15. In terms of insulin sensitivity/ glucose tolerance, studies have tended to show beneficial effects of supplementation when more sensitive measures, such as the euglycaemic hyperinsulinaemic clamp, have been used with fewer benefits seen in those using fasting measures. However, health benefits have only generally been observed in subjects at higher metabolic risk, with very little evidence to suggest that insulin sensitivity or glucose tolerance can be further improved in healthy subjects. A wide range of supplements have been used in these studies, with few direct comparisons between types of fibre, making it difficult to draw conclusions about differential health benefits.
16. Only a small number of studies have investigated the effects of isolated oligosaccharides (Abrams *et al* 2007; Brighenti *et al* 1999; Cani *et al* 2006; Castiglia-Delavaud *et al* 1998; Daubioul *et al* 2005; Ellegard *et al* 1997; Giacco *et al* 2004; Jackson *et al* 1999; Luo *et al* 1996; Schaafsma *et al* 1998; Whelan *et al* 2006), resistant starch (Noakes *et al* 1996; Park *et al* 2004; de Roos *et al* 1995; Robertson *et al* 2005) or polydextrose (Schwab *et al* 2006). None have used lignin in isolation. Studies suggest that resistant starch has a positive effect on insulin sensitivity and glucose tolerance, however, further research is required to confirm this relationship. There is no consistent evidence to support an association with weight change or maintenance of body weight in either lean or overweight subjects. There is insufficient evidence to show an association between oligosaccharides, inulin, or polydextrose on weight outcomes or metabolic profiles.

¹ Intervention studies investigating weight and metabolic outcomes are listed within the reference section.

2.2 Cardiovascular disease

17. Studies investigating fibre and cardiovascular disease have either focussed on its effect on lipids or disease endpoints such as myocardial infarction, coronary heart disease, stroke and death.

2.2.1 Observational studies

18. The cross-sectional McKeown et al 2002; Newby et al 2007; Lairon et al 2005; Ylonen et al 2003) and prospective studies (Ludwig et al 1999; Pietinen et al 1996; Wolk et al 1999; Liu et al 2000; Liu et al 2002; Jensen et al 2004) are inconclusive in their findings on the relationship between dietary fibre and risk of cardiovascular disease. Where an inverse association has been observed, it is often the association with the whole grains which have been studied. Very few studies have investigated whether there is an association with actual cardiovascular disease events. In terms of lowering cholesterol the findings are inconsistent, thus making it difficult to draw any firm conclusions.

2.2.2 Intervention Studies

19. Intervention studies (67 studies summarised in a meta-analysis by Brown et al 1999) have looked at the consequences of different types of fibre on lipid profiles. The types of fibre material studied include; oats, psyllium, pectin, barley, wheat, fibre supplements, legumes, isolated polysaccharides (Konjac-mannan, arabinogalactan, arabinoxylan, carob pulp, guar gum and hydroxypropylmethylcellulose), resistant starch and oligosaccharides and inulin. A meta analysis of intervention studies and trials subsequent to this have been considered. Overall, they suggest that soluble fibre, in particular that from oats, psyllium, pectin and guar gum may be effective in lowering total cholesterol and LDL-C when these fibres are present in the appropriate form and quantity.
20. Three intervention studies (Behall et al 2004a; Behall et al 2004b; Keenan et al 2007) that investigated barley products were identified, these suggested that fibre from this source could be effective in lowering total and LDL-C concentrations in hypercholesterolemic subjects.
21. Resistant starch does not demonstrate a cholesterol lowering effect (Heijnen et al 1996; Noakes et al 1996) and thus, based on this criterion it is not suitable to be considered a dietary fibre. There is not enough evidence to suggest that fibre from wheat, isolated polysaccharides, fibre supplement mixtures and non-digestible oligosaccharides (Arvill et al 1995; Blake et al 1997; Maki et al 2000; Vuksan et al 2000; Davidson et al 1998; Jackson et al 1999; Causey et al 2000; Luo et al 1996; Pedersen et al 1997; Schaafsma et al 1998; Brighenti et al 1999; van Dokkum et al 1999; Kruse et al 1999; Letexier et al 2003) lower cholesterol levels. In terms of legumes, three out of the four studies (Cobiac et al 1990; Fruhbeck et al 1997; Pittaway et al 2006; Anderson et al 1990) suggested that fibre from this source may have a cholesterol lowering effect, however the studies have serious flaws in study design, thus making it difficult to confirm any associations observed.
22. In terms of blood pressure, there is insufficient evidence, both from observational (Lairon et al 2005; Ascherio et al 1996) and intervention studies (24 trials included in meta

analysis by Streppel et al 2005), to demonstrate that dietary fibre has any effect on this outcome.

2.3 Colorectal cancer

2.3.1 Prospective studies

23. Cohort studies on dietary fibre and colorectal cancer generally focus on total fibre as determined by the AOAC method which does not allow separation into the different components. Therefore the individual fibre components contained in this AOAC defined fibre cannot be directly associated with risk of colorectal cancer.
24. All but one cohort study (Pietinen *et al* 1999 who used the Englyst method) assessed AOAC defined dietary fibre. In addition the length of follow up varied.
25. The evidence is variable with eight out of 17 cohort studies finding no statistically significant association between AOAC determined dietary fibre and colorectal cancer². In several studies (Willett *et al* 1990; Giovannucci *et al* 1994; Michels *et al* 2005; Schatzkin *et al* 2007; Nomura *et al* 2007) where an association was initially observed this disappeared when the studies were adjusted for multiple confounding factors. There are no reliable biomarkers of fibre intake, and it is possible that measurement error in dietary assessment leading to misclassification of exposure may have affected the results of the observational studies.
26. Ten studies have investigated the link between fruit, vegetable and grain sources of fibre and colorectal cancer (Fuchs *et al* 1999; Giovannucci *et al* 1994; Lin *et al* 2005; Mai *et al* 2003; Michels *et al* 2005; Bingham *et al* 2003; Bingham *et al* 2005; Nomura *et al* 2007; Schatzkin *et al* 2007; Terry *et al* 2001; Wakai *et al* 2007; Willett *et al.*, 1990). Whilst studies on fibre from fruit or grain generally suggest a protective effect, studies on vegetable fibre and colorectal cancer are inconclusive.
27. Three studies have distinguished between fibre from fruit, vegetable and grain sources and risk of colorectal adenoma (Giovannucci *et al* 1992; Jacobs *et al* 2002; Platz *et al* 1997), while these suggest a protective effect for fruit fibre the results for vegetable and grain fibre were inconclusive .

2.3.2 Intervention studies

28. Two RCTs (The Wheat Bran Fibre Trial (WBFT) and European Cancer Prevention Organisation Intervention Study (ECPOIS)) looked at fibre and colorectal adenoma reoccurrence (Alberts *et al* 2000; Bonithon-Kopp *et al* 2000). These both investigated wheat bran and psyllium supplements respectively. These RCTs showed either no evidence of effect when using wheat bran fibre or evidence of increased risk when using psyllium. However, the WBFT did not reach statistical significance and the ECPOIS was stopped early for financial reasons. Therefore there is insufficient evidence from trials for any of the potential dietary fibre components.

² Cohort studies investigating colorectal cancer and fibre, as measured by the AOAC method, are listed within the reference section.

2.3.3 Summary of evidence from observational and intervention studies from WCRF report

29. The working group note that the World Cancer Research Fund expert panel (WCRF 2007) were unable to draw a firm conclusion on foods containing dietary fibre (which was defined as including both foods naturally containing 'fibre' and foods which have had 'fibre' added). This was despite an apparent dose-response relationship based on cohort studies and evidence for a plausible mechanism because residual confounding could not be excluded.

2.4 Colonic function

2.4.1 Observational studies

30. Colonic function as determined by faecal weight has been demonstrated to relate to NSP intake (Davies et al 1986; Cummings et al 1992; Birkett et al 1997) but not intakes of starch or resistant starch. In particular, studies on inulin and oligosaccharides show very little effect on faecal output.

2.4.2 Intervention studies

31. Overall oligosaccharides (Alles et al 1999; Bouhnik et al 2007; Brighenti et al 1999;; Chen et al 2001; Molis et al 1996; Ten Bruggencate 2006) and inulin (Causey et al 2000; Den Hond et al 2000; Sairanen et al 2007) have very little effect on faecal weight when these materials are fed and hence an effect on faecal weight is not a criterion that is satisfied by these materials to enable them to be considered dietary fibre.
32. Overall resistant starch has a significant but modest effect on faecal weight (Jenkins et al 1998; Behall et al 2002; Muir et al 2004; Grubbens et al 2001) and this could not be considered of a size which would make it suitable to be considered dietary fibre according to this criterion.
33. The mean increase in daily faecal weight was greater for components such as wheat bran (5.4g/g) followed by fruit and vegetables (4.1g/g) and gums such as psyllium (4g/g) and least for soya products (2.5g/g) and pectin (1.2g/g) (Cummings 2001). It should be noted that many of these studies were only on a small number of subjects and were insufficiently powered

2.5 Prebiotics

34. It has been suggested that non digestible oligosaccharides such as fructooligosaccharides (FOS) , gluco-oligosaccharides (GOS) and lactulose affect the pattern of bacteria making up the gut microflora, but as yet there is no convincing evidence that as potential fibre components they confer any specific health benefit (Bouhnik et al 1996; . Bouhnik et al 1999; Bouhnik et al 2007; Gibson et al 1995; Guigoz et al 2002; Depeint et al 2008; Ito et al 1990; Ito et al 1993; Tanaka et al 1983; Tuohy et al 2002; Ballongue et al 1997).
35. Studies investigating prebiotics were of short durations (often days or weeks) and on a very small number of human subjects. The working group concluded that there was insufficient evidence from good quality human intervention studies to demonstrate that prebiotics have any health benefits which relate to their proposed actions as fibre.

References

- Abrams SA, Griffin IJ, Hawthorne KM, Liang L, Gunn SK, Darlington G & Ellis KJ (2005) A combination of prebiotic short- and long-chain inulin-type fructans enhances calcium absorption and bone mineralization in young adolescents. *American Journal of Clinical Nutrition* **82**, 471-476.
- Alberts DS, Martinez ME, Roe DJ, Guillen-Rodriguez JM, Marshall JR, van Leeuwen JB et al. (2000) Lack of effect of a high-fiber cereal supplement on the recurrence of colorectal adenomas. Phoenix Colon Cancer Prevention Physicians' Network. *N Engl J Med*; **342** (16), 1156-1162.
- Alles MS, Hartemink R, Meyboom S, Harryvan JL, Van Laere KM, Nagengast FM, Hautvast JG. (1999) Effect of transgalactooligosaccharides on the composition of the human intestinal microflora and on putative risk markers for colon cancer. *Am J Clin Nutr* **69**,980-91.
- Anderson JW, Gustafson NJ, Spencer DB, Tietyen J & Bryant CA (1990) Serum lipid response of hypercholesterolemic men to single and divided doses of canned beans *Am J Clin Nutr* **51**, 1013-9.
- Arvill A & Bodin L (1995) Effect of short-term ingestion of konjac glucomannan on serum cholesterol in healthy men *Am J Clin Nutr* **61**, 585-9.
- Ascherio A, Hennekens C, Willett WC, Sacks F, Rosner B, Manson J, Witteman J & Stampfer MJ (1996) Prospective study of nutritional factors, blood pressure, and hypertension among US women *Hypertension* **27**, 1065-72.
- Ballongue J, Schumann C & Quignon P (1997) Effects of lactulose and lactitol on colonic microflora and enzymatic activity. *Scandinavian Journal of Gastroenterology* **32**, 41-44.
- Bonithon-Kopp C, Kronborg O, Giacosa A, Rath U, Faivre J. (2000) Calcium and fibre supplementation in prevention of colorectal adenoma recurrence: a randomised intervention trial. European Cancer Prevention Organisation Study Group. *Lancet* **356** (9238), 1300-1306.
- Behall KM, Howe JC, Anderson RA. (2002) Apparent mineral retention is similar in control and hyperinsulinemic men after consumption of high amylose cornstarch. *J Nutr.* **132**:1886-91.
- Behall KM, Scholfield DJ & Hallfrisch J (2004) Diets containing barley significantly reduce lipids in mildly hypercholesterolemic men and women *Am J Clin Nutr* **80**, 1185-93.
- Behall KM, Scholfield DJ & Hallfrisch J (2004) Lipids significantly reduced by diets containing barley in moderately hypercholesterolemic men *J Am Coll Nutr* **23**, 55-62.
- Bingham SA, Day NE, Luben R, Ferrari P, Slimani N, Norat T, Clavel-Chapelon F, Kesse E, Nieters A, Boeing H, Tjønneland A, Overvad K, Martinez C, Dorransoro M, Gonzalez CA, Key TJ, Trichopoulou A, Naska A, Vineis P, Tumino R, Krogh V, Bueno-de-Mesquita HB, Peeters PH, Berglund G, Hallmans G, Lund E, Skeie G, Kaaks R, Riboli E. (2003); European Prospective Investigation into Cancer and Nutrition. Dietary fibre in food and protection against colorectal cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC): an observational study. *Lancet*.; **361** (9368):1496-501.
- Bingham SA, Norat T, Moskal A, Ferrari P, Slimani N, Clavel-Chapelon F, Kesse E, Nieters A, Boeing H, Tjønneland A, Overvad K, Martinez C, Dorransoro M, González CA, Ardanaz E, Navarro C, Quirós JR, Key TJ, Day NE, Trichopoulou A, Naska A, Krogh V, Tumino R, Palli D, Panico S, Vineis P, Bueno-de-Mesquita HB, Ocké MC, Peeters PH, Berglund G, Hallmans G, Lund E, Skeie G, Kaaks R, Riboli E. (2005); Is the association with fiber from foods in colorectal cancer confounded by folate intake? *Cancer Epidemiol Biomarkers Prev.* **14** (6):1552-6.

Birkett AM, Jones GP, de Silva AM, Young GP, Muir JG. (1997) Dietary intake and faecal excretion of carbohydrate by Australians: importance of achieving stool weights greater than 150 g to improve faecal markers relevant to colon cancer risk. *Eur J Clin Nutr.* **51**, 625-32.

Blake DE, Hamblett CJ, Frost PG, Judd PA & Ellis PR (1997) Wheat bread supplemented with depolymerized guar gum reduces the plasma cholesterol concentration in hypercholesterolemic human subjects *Am J Clin Nutr* **65**, 107-13.

Bouhnik Y, Flourie B, Riottot M, Bisetti N, Gailing M, Guibert A, Bornet F and Rambaud, JC (1996) Effects of fructooligosaccharide ingestion on fecal bifidobacteria and selected metabolic indexes of colon carcinogenesis in humans. *Nutrition and Cancer* **26**, 21-29.

Bouhnik Y, Vahedi K, Achour L, Attar A, Salfati J, Pochart P, Marteau P, Flourie B, Bornet F & Rambaud JC (1999) Short-chain fructo-oligosaccharide administration dose- dependently increases fecal bifidobacteria in healthy humans. *Journal of Nutrition* **129**, 113-116.

Bouhnik Y, Achour L, Paineau D, Riottot M, Attar A & Bornet F (2007) Four-week short chain fructo-oligosaccharides ingestion leads to increasing fecal bifidobacteria and cholesterol excretion in healthy elderly volunteers. *Nutrition Journal* **6**, 42-46.

Brighenti F, Casiraghi MC, Canzi E & Ferrari A (1999) Effect of consumption of a ready-to-eat breakfast cereal containing inulin on the intestinal milieu and blood lipids in healthy male volunteers *Eur J Clin Nutr* **53**, 726-33.

Brighenti F, Benini L, Del Rio D, Casiraghi C, Pellegrini N, Scazzina F, Jenkins DJ & Vantini I (2006) Colonic fermentation of indigestible carbohydrates contributes to the second-meal effect *Am J Clin Nutr* **83**(4), 817-22.

Brown L, Rosner B, Willett WW & Sacks FM (1999) Cholesterol-lowering effects of dietary fiber: a meta-analysis *Am J Clin Nutr* **69**, 30-42.

Causey JL, Feirtag JM, Gallaher DD, Tunland BC, Slavin JL. (2000) Effects of Dietary Inulin on Serum Lipids, Blood Glucose and the Gastrointestinal Environment in Hypercholesterolemic men. *Nutrition Res* **20**, 191-201.

Cani PD, Joly E, Horsmans Y & Delzenne NM (2006) Oligofructose promotes satiety in healthy human: a pilot study *Eur J Clin Nutr* **60**(5), 567-72.

Castiglia-Delavaud C, Verdier E, Besle JM, Vernet J, Boirie Y, Beaufriere B, De Baynast R, Vermorel M. (1998) Net energy value of non-starch polysaccharide isolates (sugarbeet fibre and commercial inulin) and their impact on nutrientdigestive utilization in healthy human subjects. *Br J Nutr.* **80**, 343-52.

Colditz GA, Manson JE, Stampfer MJ, Rosner B, Willett WC & Speizer FE (1992) Diet and risk of clinical diabetes in women *Am J Clin Nutr* **55**(5), 1018-23.

Chen HL, Lu YH, Lin JJ, Ko LY. (2001) Effects of isomalto-oligosaccharides on bowel functions and indicators of nutritional status in constipated elderly men. *J Am Coll Nutr.* **20**, 44-9.

Cobiac L, McArthur R & Nestel PJ (1990) Can eating baked beans lower plasma cholesterol? *Eur J Clin Nutr* **44**, 819-22.

Cummings JH, Bingham SA, Heaton KW, Eastwood MA. (1992) Fecal weight, colon cancer risk, and dietary intake of nonstarch polysaccharides (dietary fiber). *Gastroenterology.* **103**, 1783-9.

Cummings JH. (2001) The effect of dietary fiber on fecal weight and composition. Chapter 4.4 in: *CRC Handbook of Dietary Fiber in Human Nutrition*. 3rd edition. Edit GA Spiller. *CRC Press*, Boca Raton.

Davies GJ, Crowder M, Reid B, Dickerson JW. (1986) Bowel function measurements of individuals with different eating patterns. *Gut.* **27**, 164-9.

Davidson MH, Maki KC, Kong JC, Dugan LD, Torri SA, Hall HA, Drennan KB, Anderson SM, Fulgoni VL, Saldanha LG & Olson BH (1998) Long-term effects of consuming foods containing psyllium seed husk on serum lipids in subjects with hypercholesterolemia *Am J Clin Nutr* **67**, 367-76.

Den Hond E, Geypens B, Ghoois Y. (2000) Effect of High Performance Chicory Inulin on Constipation. *Nutrition Res* **20**, 731-36.

Depeint F, Tzortzis G, Vulevic J, Tanson, K & Gibson GR (2008) Prebiotic evaluation of a novel galactooligosaccharide mixture produced by the enzymatic activity of *Bifidobacterium bifidum* NCIMB 41171, in healthy humans: a randomised, double-blind, crossover, placebo-controlled intervention study. *American Journal of Clinical Nutrition* **87**, 785-791.

Department of Health. (1991). Dietary Reference Values for Food Energy and Nutrients for the United Kingdom. Report of the panel of Dietary Reference Values of the Committee on Medical Aspects of Food Policy. Report on Health and Social Subjects **41**. HMSO: London.

de Roos N, Heijnen ML, de Graaf C, Woestenenk G & Hobbel E (1995) Resistant starch has little effect on appetite, food intake and insulin secretion of healthy young men *Eur J Clin Nutr* **49**(7), 532-41.

Ellegård L, Andersson H & Bosaeus I (1997) Inulin and oligofructose do not influence the absorption of cholesterol, or the excretion of cholesterol, Ca, Mg, Zn, Fe or bile acids but increases energy excretion in ileostomy subjects. *European Journal of Clinical Nutrition* **51**, 1-5.

Fruhbeck G, Monreal I & Santidrian S (1997) Hormonal implications of the hypocholesterolemic effect of intake of field beans (*Vicia faba* L.) by young men with hypercholesterolemia *Am J Clin Nutr* **66**, 1452-60.

Giacco R, Clemente G, Luongo D, Lasorella G, Fiume I, Brouns F, Bornet F, Patti L, Cipriano P, Rivellese AA & Riccardi G (2004) Effects of short-chain fructooligosaccharides on glucose and lipid metabolism in mild hypercholesterolaemic individuals *Clin Nutr* **23**(3), 331-40.

Gibson GR & Roberfroid MB (1995) Dietary modulation of the human colonic microflora introducing the concept of probiotics. *Journal of Nutrition* **125**, 1401-1412.

Giovannucci E, Rimm EB, Stampfer MJ, Colditz GA, Ascherio A, Willett WC (1994) Intake of fat, meat, and fiber in relation to risk of colon cancer in men. *Cancer Res* **54**(9):2390-2397.

Giovannucci E, Stampfer MJ, Colditz G, Rimm EB, Willett WC. (1992) Relationship of diet to risk of colorectal adenoma in men. *J Natl Cancer Inst* **84**(2), 91-98.

Grubben MJ, van den Braak CC, Essenberg M, Olthof M, Tangerman A, Katan MB, Nagengast FM. (2001) Effect of resistant starch on potential biomarkers for colonic cancer risk in patients with colonic adenomas: a controlled trial. *Dig Dis Sci*. **46**, 750-6.

Guigoz Y, Rochat F, Perruisseau-Carrier G, Rochat I & Schriffin EJ (2002) Effects of oligosaccharide on the fecal flora and non-specific immune system in elderly people. *Nutrition Reviews* **22**, 13-25.

Heijnen ML, van Amelsvoort JM, Deurenberg P & Beynen AC (1996) Neither raw nor retrograded resistant starch lowers fasting serum cholesterol concentrations in healthy normolipidemic subjects *Am J Clin Nutr* **64**, 312-8.

Hodge AM, English DR, O'Dea K & Giles GG (2004) Glycemic index and dietary fiber and the risk of type 2 diabetes *Diabetes Care* **27**(11), 2701-6.

Hu FB, Manson JE, Stampfer MJ, Colditz G, Liu S, Solomon CG & Willett WC (2001) Diet, lifestyle, and the risk of type 2 diabetes mellitus in women *N Engl J Med* **345**(11), 790-7.

- Iqbal SI, Helge JW & Heitmann BL (2006) Do energy density and dietary fiber influence subsequent 5-year weight changes in adult men and women? *Obesity* (Silver Spring) **14**(1), 106-14.
- Ito M, Deguchi Y, Miyamori A, Matsumoto K, Kikuchi H, Matsumoto K, Kobayashi Y, Yajima T & Kan T (1990) Effects of administration of galactooligosaccharides on the human fecal microflora, stool weight and abdominal sensation. *Microbial Ecology in Health and Disease* **3**, 285-292.
- Ito M, Kimura M, Deguchi Y, Miyamori-Watabe A, Yajima T & Kan T (1993) Effects of *trans*galactosylated disaccharides on the human intestinal microflora and their metabolism. *Journal of Nutritional Science and Vitaminology* **39**, 279-288.
- Jacobs ET, Giuliano AR, Roe DJ, Guillen-Rodriguez JM, Hess LM, Alberts DS et al. (2002); Intake of supplemental and total fiber and risk of colorectal adenoma recurrence in the wheat bran fiber trial. *Cancer Epidemiol Biomarkers Prev* **11** (9), 906-914.
- Jackson KG, Taylor GR, Clohessy AM & Williams CM (1999) The effect of the daily intake of inulin on fasting lipid, insulin and glucose concentrations in middle-aged men and women *Br J Nutr* **82**, 23-30.
- Jenkins DJ, Vuksan V, Kendall CW, Würsch P, Jeffcoat R, Waring S, Mehling CC, Vidgen E, Augustin LS, Wong E. (1998) Physiological effects of resistant starches on fecal bulk, short chain fatty acids, blood lipids and glycemic index. *J Am Coll Nutr.* **17**, 609-16.
- Jensen MK, Koh-Banerjee P, Hu FB, Franz M, Sampson L, Gronbaek M & Rimm EB (2004) Intakes of whole grains, bran, and germ and the risk of coronary heart disease in men *Am J Clin Nutr* **80**, 1492-9.
- Keenan JM, Goulson M, Shamliyan T, Knutson N, Kolberg L & Curry L (2007) The effects of concentrated barley beta-glucan on blood lipids in a population of hypercholesterolaemic men and women *Br J Nutr* **97**, 1162-8.
- Krishnan S, Rosenberg L, Singer M, Hu FB, Djousse L, Cupples LA & Palmer JR (2007) Glycemic index, glycemic load, and cereal fiber intake and risk of type 2 diabetes in US black women *Arch Intern Med* **167**(21), 2304-9.
- Kruse HP, Kleessen B & Blaut M (1999) Effects of inulin on faecal bifidobacteria in human subjects. *British Journal of Nutrition* **82**, 375-382.
- Lairon D, Arnault N, Bertrais S, Planells R, Clero E, Hercberg S & Boutron-Ruault MC (2005) Dietary fiber intake and risk factors for cardiovascular disease in French adults *Am J Clin Nutr* **82**, 1185-94.
- Letexier D, Diraison F & Beylot M (2003) Addition of inulin to a moderately high-carbohydrate diet reduces hepatic lipogenesis and plasma triacylglycerol concentrations in humans *Am J Clin Nutr* **77**, 559-64.
- Lindstrom J, Ilanne-Parikka P, Peltonen M, Aunola S, Eriksson JG, Hemio K, Hamalainen H, Harkonen P, Keinanen-Kiukaanniemi S, Laakso M, Louheranta A, Mannelin M, Paturi M, Sundvall J, Valle TT, Uusitupa M & Tuomilehto J (2006) Sustained reduction in the incidence of type 2 diabetes by lifestyle intervention: follow-up of the Finnish Diabetes Prevention Study *Lancet* **368**(9548), 1673-9.
- Lindstrom J, Peltonen M, Eriksson JG, Louheranta A, Fogelholm M, Uusitupa M & Tuomilehto J (2006) High-fibre, low-fat diet predicts long-term weight loss and decreased type 2 diabetes risk: the Finnish Diabetes Prevention Study *Diabetologia* **49**(5), 912-20.
- Liu S, Buring JE, Sesso HD, Rimm EB, Willett WC & Manson JE (2002) A prospective study of dietary fiber intake and risk of cardiovascular disease among women *J Am Coll Cardiol* **39**, 49-56.
- Ludwig DS, Pereira MA, Kroenke CH, Hilner JE, Van Horn L, Slattery ML & Jacobs DR, Jr. (1999) Dietary fiber, weight gain, and cardiovascular disease risk factors in young adults *Jama* **282**, 1539-46.

- Luo J, Rizkalla SW, Alamowitch C, Boussairi A, Blayo A, Barry JL, Laffitte A, Guyon F, Bornet FR & Slama G (1996) Chronic consumption of short-chain fructooligosaccharides by healthy subjects decreased basal hepatic glucose production but had no effect on insulin-stimulated glucose metabolism *Am J Clin Nutr* **63**, 939-45.
- Maki KC, Davidson MH, Torri S, Ingram KA, O'Mullane J, Daggy BP & Albrecht HH (2000) High-molecular-weight hydroxypropylmethylcellulose taken with or between meals is hypocholesterolemic in adult men *J Nutr* **130**, 1705-10.
- McKeown NM, Meigs JB, Liu S, Wilson PW & Jacques PF (2002) Whole-grain intake is favorably associated with metabolic risk factors for type 2 diabetes and cardiovascular disease in the Framingham Offspring Study *Am J Clin Nutr* **76**, 390-8.
- Meyer KA, Kushi LH, Jacobs DR, Jr., Slavin J, Sellers TA & Folsom AR (2000) Carbohydrates, dietary fiber, and incident type 2 diabetes in older women *Am J Clin Nutr* **71**(4), 921-30.
- Michels KB, Fuchs CS, Giovannucci E, Colditz GA, Hunter DJ, Stampfer MJ et al. (2005) Fiber intake and incidence of colorectal cancer among 76,947 women and 47,279 men. *Cancer Epidemiol Biomarkers Prev* **14** (4):842-849.
- Molis C, Flourié B, Ouarné F, Gailing MF, Lartigue S, Guibert A, Bornet F, Galmiche JP. (1996) Digestion, excretion, and energy value of fructooligosaccharides in healthy humans. *Am J Clin Nutr.* **64**, 324-8.
- Montonen J, Knekt P, Jarvinen R, Aromaa A & Reunanen A (2003) Whole-grain and fiber intake and the incidence of type 2 diabetes *Am J Clin Nutr* **77**(3), 622-9.
- Newby PK, Maras J, Bakun P, Muller D, Ferrucci L & Tucker KL (2007) Intake of whole grains, refined grains, and cereal fiber measured with 7-d diet records and associations with risk factors for chronic disease *Am J Clin Nutr* **86**, 1745-53.
- Muir JG, Yeow EG, Keogh J, Pizzey C, Bird AR, Sharpe K, O'Dea K, Macrae FA. (2004) Combining wheat bran with resistant starch has more beneficial effects on fecal indexes than does wheat bran alone. *Am J Clin Nutr.* **79**, 1020-8.
- Newby PK, Peterson KE, Berkey CS, Leppert J, Willett WC & Colditz GA (2003) Dietary composition and weight change among low-income preschool children *Arch Pediatr Adolesc Med* **157**(8), 759-64.
- Noakes M, Clifton PM, Nestel PJ, Le Leu R & McIntosh G (1996) Effect of high-amylose starch and oat bran on metabolic variables and bowel function in subjects with hypertriglyceridemia *Am J Clin Nutr* **64**, 944-51.
- Nomura AM, Hankin JH, Henderson BE, Wilkens LR, Murphy SP, Pike MC et al. (2007) Dietary fiber and colorectal cancer risk: the multiethnic cohort study. *Cancer Causes Control* **18** (7), 753-764.
- Park OJ, Kang NE, Chang MJ & Kim WK (2004) Resistant starch supplementation influences blood lipid concentrations and glucose control in overweight subjects *J Nutr Sci Vitaminol (Tokyo)* **50**(2), 93-9.
- Pedersen A, Sandstrom B & Van Amelsvoort JM (1997) The effect of ingestion of inulin on blood lipids and gastrointestinal symptoms in healthy females *Br J Nutr* **78**, 215-22.
- Pietinen P, Rimm EB, Korhonen P, Hartman AM, Willett WC, Albanes D & Virtamo J (1996) Intake of dietary fiber and risk of coronary heart disease in a cohort of Finnish men. The Alpha-Tocopherol, Beta-Carotene Cancer Prevention Study *Circulation* **94**, 2720-7.
- Pietinen P, Malila N, Virtanen M, Hartman TJ, Tangrea JA, Albanes D et al. (1999) Diet and risk of colorectal cancer in a cohort of Finnish men. *Cancer Causes Control* **10** (5), 387-396.

Pittaway JK, Ahuja KD, Cehun M, Chronopoulos A, Robertson IK, Nestel PJ & Ball MJ (2006) Dietary supplementation with chickpeas for at least 5 weeks results in small but significant reductions in serum total and low-density lipoprotein cholesterol in adult women and men *Ann Nutr Metab* **50**, 512-8.

Platz EA, Giovannucci E, Rimm EB, Rockett HR, Stampfer MJ, Colditz GA et al. (1997) Dietary fiber and distal colorectal adenoma in men. *Cancer Epidemiol Biomarkers Prev* **6** (9), 661-670.

Robertson DJ, Sandler RS, Haile R, Tosteson TD, Greenberg ER, Grau M et al. (2005); Fat, fiber, meat and the risk of colorectal adenomas. *Am J Gastroenterol* **100** (12):2789-2795.

SACN frame work (2002) www.sacn.gov.uk

Sairanen U, Pirainen L, Gråsten S, Tompuri T, Mättö J, Saarela M, Korpela R. (2007) The effect of probiotic fermented milk and inulin on the functions and microecology of the intestine. *J Dairy Res.* **74**, 367-73.

Salmeron J, Ascherio A, Rimm EB, Colditz GA, Spiegelman D, Jenkins DJ, Stampfer MJ, Wing AL & Willett WC (1997) Dietary fiber, glycemic load, and risk of NIDDM in men *Diabetes Care* **20**(4), 545-50.

Salmeron J, Manson JE, Stampfer MJ, Colditz GA, Wing AL & Willett WC (1997) Dietary fiber, glycemic load, and risk of non-insulin-dependent diabetes mellitus in women *Jama* **277**(6), 472-7.

Schatzkin A, Mouw T, Park Y, Subar AF, Kipnis V, Hollenbeck A et al. (2007) Dietary fiber and whole-grain consumption in relation to colorectal cancer in the NIH-AARP Diet and Health Study. *Am J Clin Nutr* **85** (5), 1353-1360.

Schaafsma G, Meuling WJ, van Dokkum W & Bouley C (1998) Effects of a milk product, fermented by *Lactobacillus acidophilus* and with fructo-oligosaccharides added, on blood lipids in male volunteers *Eur J Clin Nutr* **52**, 436-40.

Schulze MB, Liu S, Rimm EB, Manson JE, Willett WC & Hu FB (2004) Glycemic index, glycemic load, and dietary fiber intake and incidence of type 2 diabetes in younger and middle-aged women *Am J Clin Nutr* **80**(2), 348-56. 190

Schulze MB, Schulz M, Heidemann C, Schienkiewitz A, Hoffmann K & Boeing H(2007) Fiber and magnesium intake and incidence of type 2 diabetes: a prospective study and meta-analysis *Arch Intern Med* **167**(9), 956-65.

Schwab U, Louheranta A, Torronen A & Uusitupa M (2006) Impact of sugar beet pectin and polydextrose on fasting and postprandial glycemia and fasting concentrations of serum total and lipoprotein lipids in middle-aged subjects with abnormal glucose metabolism *Eur J Clin Nutr* **60**, 1073-80.

Stevens J, Ahn K, Juhaeri, Houston D, Steffan L & Couper D (2002) Dietary fiber intake and glycemic index and incidence of diabetes in African-American and white adults: the ARIC study *Diabetes Care* **25**(10), 1715-21.

Streppel MT, Arends LR, van 't Veer P, Grobbee DE & Geleijnse JM (2005) Dietary fiber and blood pressure: a meta-analysis of randomised placebo-controlled trials *Arch Intern Med* **165**, 150-6.

Tanaka R, Takayama H, Morotomi M, Kuroshima T, Ueyama S, Matsumoto K, Kuroda A & Mutai M (1983) Effects of administration of TOS and *Bifidobacterium breve* 4006 on the human fecal flora. *Bifidobacteria Microflora* **2**, 17-24.

Ten Bruggencate SJ, Bovee-Oudenhoven IM, Lettink-Wissink ML, Katan MB, van der Meer R. (2006) Dietary fructooligosaccharides affect intestinal barrier function in healthy men. *J Nutr.* **136**, 70-4.

Tuohy KM, Ziemer CJ, Klinder A, Knobel Y, Pool-Zobel BL & Gibson GR (2002) A human volunteer study to determine the prebiotic effects of lactulose powder on human colonic bacteria. *Microbial Ecology in Health and Disease* **14**,165-173.

van Dokkum W, Wezendonk B, Srikumar TS & van den Heuvel EG (1999) Effect of nondigestible oligosaccharides on large-bowel functions, blood lipid concentrations and glucose absorption in young healthy male subjects *Eur J Clin Nutr* **53**(1), 1-7.

Vuksan V, Sievenpiper JL, Owen R, Swilley JA, Spadafora P, Jenkins DJ, Vidgen E, Brighenti F, Josse RG, Leiter LA, Xu Z & Novokmet R (2000) Beneficial effects of viscous dietary fiber from Konjac-mannan in subjects with the insulin resistance syndrome: results of a controlled metabolic trial *Diabetes Care* **23**(1), 9-14.

Whelan K, Efthymiou L, Judd PA, Preedy VR & Taylor MA (2006) Appetite during consumption of enteral formula as a sole source of nutrition: the effect of supplementing pea-fibre and fructo-oligosaccharides *Br J Nutr* **96**(2), 350-6.

Willett WC, Stampfer MJ, Colditz GA, Rosner BA, Speizer FE. (1990); Relation of meat, fat, and fiber intake to the risk of colon cancer in a prospective study among women. *N Engl J Med* **323** (24),1664-1672.

Wolk A, Manson JE, Stampfer MJ, Colditz GA, Hu FB, Speizer FE, Hennekens CH & Willett WC (1999) Long-term intake of dietary fiber and decreased risk of coronary heart disease among women *Jama* **281**, 1998-2004.

World Cancer Research Fund/ American Institute of Cancer Research. (2007) Food Nutrition, Physical Activity, and the Prevention of Cancer: a Global Perspective...Washington DC: AICR.

Intervention studies investigating weight and obesity

Astrup A, Vrist E & Quaade F (1990) Dietary fibre added to very low calorie diet reduces hunger and alleviates constipation *Int J Obes* **14**(2), 105-12.

Barroso Aranda J, Contreras F, Bagchi D & Preuss HG (2002) Efficacy of a novel chitosan formulation on fecal fat excretion: a double-blind, crossover, placebocontrolled study *J Med* **33**(1-4), 209-25.

Birketvedt GS, Aaseth J, Florholmen JR & Ryttig K (2000) Long-term effect of fibre supplement and reduced energy intake on body weight and blood lipids in overweight subjects *Acta Medica (Hradec Kralove)* **43**(4), 129-32.

Birketvedt GS, Shimshi M, Erling T & Florholmen J (2005) Experiences with three different fiber supplements in weight reduction *Med Sci Monit* **11**(1), PI5-8.

Cairella G, Cairella M & Marchini G (1995) Effect of dietary fibre on weight correction after modified fasting *Eur J Clin Nutr* **49** Suppl 3, S325-7.

Cicero AF, Derosa G, Manca M, Bove M, Borghi C & Gaddi AV (2007) Different effect of psyllium and guar dietary supplementation on blood pressure control in hypertensive overweight patients: a six-month, randomised clinical trial *Clin Exp Hypertens* **29**(6), 383-94.

Davy BM, Davy KP, Ho RC, Beske SD, Davrath LR & Melby CL (2002) High-fiber oat cereal compared with wheat cereal consumption favorably alters LDL-cholesterol subclass and particle numbers in middle-aged and older men *Am J Clin Nutr* **76**, 351-8.

Duncan LJ, Rose K & Meiklejohn AP (1960) Phenmetrazine hydrochloride and methylcellulose in the treatment of "refractory" obesity *Lancet* **1**(7137), 1262-5.

Effertz ME, Denman P & Slavin JL (1991) The effect of soy polysaccharide on body weight, serum lipids, blood glucose and fecal parameters in moderately obese adults *Nutrition Research* **11**(8), 849-859.

Eliasson K, Ryttig KR, Hylander B & Rossner S (1992) A dietary fibre supplement in the treatment of mild hypertension. A randomised, double-blind placebo-controlled trial *J Hypertens* **10**(2), 195-9.

Evans E & Miller DS (1975) Bulking agents in the treatment of obesity *Nutr Metab* **18**(4), 199-203.

- Frost GS, Brynes AE, Dhillon WS, Bloom SR & McBurney MI (2003) The effects of fiber enrichment of pasta and fat content on gastric emptying, GLP-1, glucose, and insulin responses to a meal *Eur J Clin Nutr* **57**(2), 293-8.
- Howard BV, Manson JE, Stefanick ML, Beresford SA, Frank G, Jones B, Rodabough RJ, Snetselaar L, Thomson C, Tinker L, Vitolins M & Prentice R (2006) Low-fat dietary pattern and weight change over 7 years: the Women's Health Initiative Dietary Modification Trial *Jama* **295**(1), 39-49.
- Howarth NC, Saltzman E, McCrory MA, Greenberg AS, Dwyer J, Ausman L, Kramer DG & Roberts SB (2003) Fermentable and nonfermentable fiber supplements did not alter hunger, satiety or body weight in a pilot study of men and women consuming self-selected diets *J Nutr* **133**(10), 3141-4.
- Hylander B & Rossner S (1983) Effects of dietary fiber intake before meals on weight loss and hunger in a weight-reducing club *Acta Med Scand* **213**(3), 217-20.
- Jenkins DJ, Wolever TM, Rao AV, Hegele RA, Mitchell SJ, Ransom TP, Boctor DL, Spadafora PJ, Jenkins AL, Mehling C & et al. (1993) Effect on blood lipids of very high intakes of fiber in diets low in saturated fat and cholesterol *N Engl J Med* **329**(1), 21-6.
- Kovacs EM, Westerterp-Plantenga MS, Saris WH, Goossens I, Geurten P & Brouns F (2001) The effect of addition of modified guar gum to a low-energy semisolid meal on appetite and body weight loss *Int J Obes Relat Metab Disord* **25**(3), 307-15.
- Lo GS & Cole TG (1990) Soy cotyledon fiber products reduce plasma lipids *Atherosclerosis* **82**(1-2), 59-67.
- Mackay S & Ball MJ (1992) Do beans and oat bran add to the effectiveness of a low-fat diet? *Eur J Clin Nutr* **46**(9), 641-8.
- Marett R & Slavin JL (2004) No long-term benefits of supplementation with arabinogalactan on serum lipids and glucose *J Am Diet Assoc* **104**, 636-9.
- Pasman WJ, Saris WH, Wauters MA & Westerterp-Plantenga MS (1997) Effect of one week of fibre supplementation on hunger and satiety ratings and energy intake *Appetite* **29**(1), 77-87.
- Pasman WJ, Westerterp-Plantenga MS, Muls E, Vansant G, van Ree J & Saris WH (1997) The effectiveness of long-term fibre supplementation on weight maintenance in weight-reduced women *Int J Obes Relat Metab Disord* **21**(7), 548-55.
- Pelkman CL, Navia JL, Miller AE & Pohle RJ (2007) Novel calcium-gelled, alginatepectin beverage reduced energy intake in nondieting overweight and obese women: interactions with dietary restraint status *Am J Clin Nutr* **86**(6), 1595-602.
- Reyna-Villasmil N, Bermudez-Pirela V, Mengual-Moreno E, Arias N, Cano-Ponce C, Leal-Gonzalez E, Souki A, Inglett GE, Israili ZH, Hernandez-Hernandez R, Valasco M & Arraiz N (2007) Oat-derived beta-glucan significantly improves HDLC and diminishes LDLC and non-HDL cholesterol in overweight individuals with mild hypercholesterolemia *Am J Ther* **14**, 203-12.
- Rigaud D, Ryttig KR, Leeds AR, Bard D & Apfelbaum M (1987) Effects of a moderate dietary fibre supplement on hunger rating, energy input and faecal energy output in young, healthy volunteers. A randomised, double-blind, cross-over trial *Int J Obes* **11** Suppl 1, 73-8.
- Rigaud D, Ryttig KR, Angel LA & Apfelbaum M (1990) Overweight treated with energy restriction and a dietary fibre supplement: a 6-month randomised, doubleblind, placebo-controlled trial *Int J Obes* **14**(9), 763-9.
- Ross AH, Eastwood MA, Brydon WG, Anderson JR & Anderson DM (1983) A study of the effects of dietary gum arabic in humans *Am J Clin Nutr* **37**(3), 368-75.

- Rossner S, Andersson IL & Ryttig K (1988) Effects of a dietary fibre supplement to a weight reduction programme on blood pressure. A randomised, double-blind, placebo-controlled study *Acta Med Scand* **223**(4), 353-7.
- Rossner S, von Zweigbergk D, Ohlin A & Ryttig K (1987) Weight reduction with dietary fibre supplements. Results of two double-blind randomised studies *Acta Med Scand* **222**(1), 83-8.
- Ryttig KR, Lammert O, Nielsen E, Garby L & Poulsen K (1990) The effect of a soluble dietary fibre supplement on 24-hour energy expenditure during a standardized physical activity programme *Int J Obes* **14**(5), 451-5.
- Ryttig KR, Tellnes G, Haegh L, Boe E & Fagerthun H (1989) A dietary fibre supplement and weight maintenance after weight reduction: a randomised, double-blind, placebo-controlled long-term trial *Int J Obes* **13**(2), 165-71.
- Saltzman E, Moriguti JC, Das SK, Corrales A, Fuss P, Greenberg AS & Roberts SB (2001) Effects of a cereal rich in soluble fiber on body composition and dietary compliance during consumption of a hypocaloric diet *J Am Coll Nutr* **20**(1), 50-7.
- Solum TT, Ryttig KR, Solum E & Larsen S (1987) The influence of a high-fibre diet on body weight, serum lipids and blood pressure in slightly overweight persons. A randomised, double-blind, placebo-controlled investigation with diet and fibre tablets (DumoVital) *Int J Obes* **11** Suppl 1, 67-71.
- Southgate DA, Branch WJ, Hill MJ, Drasar BS, Walters RL, Davies PS & Baird IM (1976) Metabolic responses to dietary supplements of bran *Metabolism* **25**(10), 1129-35.
- Stevens J, Levitsky DA, VanSoest PJ, Robertson JB, Kalkwarf HJ & Roe DA (1987) Effect of psyllium gum and wheat bran on spontaneous energy intake *Am J Clin Nutr* **46**(5), 812-7.
- Tuomilehto J, Voutilainen E, Huttunen J, Vinni S & Homan K (1980) Effect of guar gum on body weight and serum lipids in hypercholesterolemic females *Acta Med Scand* **208**(1-2), 45-8.
- Vajifdar BU, Goyal VS, Lokhandwala YY, Mhamunkar SR, Mahadik SP, Gawad AK, Halankar SA & Kulkarni HL (2000) Is dietary fiber beneficial in chronic ischemic heart disease? *J Assoc Physicians India* **48**(9), 871-6.
- Valle-Jones JC (1980) The evaluation of a new appetite-reducing agent (Prefil) in the management of obesity *Br J Clin Pract* **34**(3), 72-4.
- Walsh DE, Yaghoubian V & Behforooz A (1984) Effect of glucomannan on obese patients: a clinical study *Int J Obes* **8**(4), 289-93.
- Wisker E, Maltz A & Feldheim W (1988) Metabolizable energy of diets low or high in dietary fiber from cereals when eaten by humans *J Nutr* **118**(8), 945-52.
- Yudkin J (1959) The causes and cure of obesity *Lancet* **2**(7112), 1135-8.

Intervention studies investigating metabolic risk factors

- Aller R, de Luis DA, Izaola O, La Calle F, del Olmo L, Fernandez L, Arranz T & Hernandez JM (2004) Effect of soluble fiber intake in lipid and glucose levels in healthy subjects: a randomised clinical trial *Diabetes Res Clin Pract* **65**, 7-11.
- Chearskul S, Supingklud N, Nitithamyong A & Sirichakwal P (2006) Assessment of hormonal and metabolic effects of dietary fiber in young Thai women *J Med Assoc Thai* **89**(7), 997-1003.
- Garcia AL, Otto B, Reich SC, Weickert MO, Steiniger J, Machowetz A, Rudovich NN, Mohlig M, Katz N, Speth M, Meuser F, Doerfer J, Zunft HJ, Pfeiffer AH & Koebnick C (2007) Arabinoxylan consumption decreases postprandial serum glucose, serum insulin and plasma total ghrelin response in subjects with impaired glucose tolerance *Eur J Clin Nutr* **61**(3), 334-41.

Hanai H, Ikuma M, Sato Y, Iida T, Hosoda Y, Matsushita I, Nogaki A, Yamada M & Kaneko E (1997) Long-term effects of water-soluble corn bran hemicellulose on glucose tolerance in obese and non-obese patients: improved insulin sensitivity and glucose metabolism in obese subjects *Biosci Biotechnol Biochem* **61**(8), 1358-61.

Jenkins DJ, Leeds AR, Gassull MA, Cochet B & Alberti GM (1977) Decrease in postprandial insulin and glucose concentrations by guar and pectin *Ann Intern Med* **86**(1), 20-3.

Keogh GF, Cooper GJ, Mulvey TB, McArdle BH, Coles GD, Monro JA & Poppitt SD (2003) Randomised controlled crossover study of the effect of a highly beta-glucan-enriched barley on cardiovascular disease risk factors in mildly hypercholesterolemic men *Am J Clin Nutr* **78**(4), 711-8.

Kestin M, Moss R, Clifton PM & Nestel PJ (1990) Comparative effects of three cereal brans on plasma lipids, blood pressure, and glucose metabolism in mildly hypercholesterolemic men *Am J Clin Nutr* **52**(4), 661-6.

Landin K, Holm G, Tengborn L & Smith U (1992) Guar gum improves insulin sensitivity, blood lipids, blood pressure, and fibrinolysis in healthy men *Am J Clin Nutr* **56**(6), 1061-5.

Li J, Kaneko T, Qin LQ, Wang J & Wang Y (2003) Effects of barley intake on glucose tolerance, lipid metabolism, and bowel function in women *Nutrition* **19**(11-12), 926-9.

Maki KC, Galant R, Samuel P, Tesser J, Witchger MS, Ribaya-Mercado JD, Blumberg JB & Geohas J (2007) Effects of consuming foods containing oat beta-glucan on blood pressure, carbohydrate metabolism and biomarkers of oxidative stress in men and women with elevated blood pressure *Eur J Clin Nutr* **61**(6), 786-95.

Munoz, J. M. & Sandstead, H. H. & Jacob, R. A. (1979) Effects of dietary fiber on glucose tolerance of normal men *Diabetes* **25**, 496-502.

Pereira MA, Jacobs DR, Jr., Pins JJ, Raatz SK, Gross MD, Slavin JL & Seaquist ER (2002) Effect of whole grains on insulin sensitivity in overweight hyperinsulinemic adults *Am J Clin Nutr* **75**(5), 848-55.

Sabovic M, Lavre S & Keber I (2004) Supplementation of wheat fibre can improve risk profile in patients with dysmetabolic cardiovascular syndrome *Eur J Cardiovasc Prev Rehabil* **11**(2), 144-8.

Vuksan V, Sievenpiper JL, Owen R, Swilley JA, Spadafora P, Jenkins DJ, Vidgen E, Brighenti F, Josse RG, Leiter LA, Xu Z & Novokmet R (2000) Beneficial effects of viscous dietary fiber from Konjac-mannan in subjects with the insulin resistance syndrome: results of a controlled metabolic trial *Diabetes Care* **23**(1), 9-14.

Weickert MO, Mohlig M, Schofl C, Arafat AM, Otto B, Viehoff H, Koebnick C, Kohl A, Spranger J & Pfeiffer AF (2006) Cereal fiber improves whole-body insulin sensitivity in overweight and obese women *Diabetes Care* **29**(4), 775-80.

Prospective studies investigating colorectal cancer (AOAC method of analysis)

Fuchs CS, Giovannucci EL, Colditz GA, Hunter DJ, Stampfer MJ, Rosner B et al. (1999) Dietary fiber and the risk of colorectal cancer and adenoma in women. *N Engl J Med* **340** (3), 169-176.

Gaard M, Tretli S, Loken EB. (1996) Dietary factors and risk of colon cancer: a prospective study of 50,535 young Norwegian men and women. *Eur J Cancer Prev* **5** (6), 445-454.

Giovannucci E, Rimm EB, Stampfer MJ, Colditz GA, Ascherio A, Willett WC (1994) Intake of fat, meat, and fiber in relation to risk of colon cancer in men. *Cancer Res* **54**(9):2390-2397.

Heilbrun LK, Nomura A, Hankin JH, Stemmermann GN. (1989) Diet and Colorectal- Cancer with Special Reference to Fiber Intake. *International Journal of Cancer* **44**(1):1-6.

Higginbotham S, Zhang ZF, Lee IM, Cook NR, Giovannucci E, Buring JE et al. (2004) Dietary glycemic load and risk of colorectal cancer in the Women's Health Study. *J Natl Cancer Inst* **96** (3), 229-233.

- Lin J, Zhang SM, Cook NR, Rexrode KM, Liu S, Manson JE et al. (2005); Dietary intakes of fruit, vegetables, and fiber, and risk of colorectal cancer in a prospective cohort of women (United States). *Cancer Causes Control* **16** (3), 225-233.
- Mai V, Flood A, Peters U, Lacey JV, Jr., Schairer C, Schatzkin A. (2003) Dietary fibre and risk of colorectal cancer in the Breast Cancer Detection Demonstration Project (BCDDP) follow-up cohort. *Int J Epidemiol* **32** (2), 234-239.
- McCullough ML, Robertson AS, Chao A, Jacobs EJ, Stampfer MJ, Jacobs DR et al. (2003) A prospective study of whole grains, fruits, vegetables and colon cancer risk. *Cancer Causes Control* **14** (10), 959-970.
- Michels KB, Fuchs CS, Giovannucci E, Colditz GA, Hunter DJ, Stampfer MJ et al. (2005) Fiber intake and incidence of colorectal cancer among 76,947 women and 47,279 men. *Cancer Epidemiol Biomarkers Prev* **14** (4):842-849.
- Nomura AM, Hankin JH, Henderson BE, Wilkens LR, Murphy SP, Pike MC et al. (2007) Dietary fiber and colorectal cancer risk: the multiethnic cohort study. *Cancer Causes Control* **18** (7), 753-764.
- Otani T, Iwasaki M, Ishihara J, Sasazuki S, Inoue M, Tsugane S. (2006) Dietary fiber intake and subsequent risk of colorectal cancer: the Japan Public Health Center-based prospective study. *Int J Cancer* **119** (6), 1475-1480.
- Schatzkin A, Mouw T, Park Y, Subar AF, Kipnis V, Hollenbeck A et al. (2007) Dietary fiber and whole-grain consumption in relation to colorectal cancer in the NIH-AARP Diet and Health Study. *Am J Clin Nutr* **85** (5), 1353-1360.
- Sellers TA, Bazyk AE, Bostick RM, Kushi LH, Olson JE, Anderson KE et al. (1998) Diet and risk of colon cancer in a large prospective study of older women: an analysis stratified on family history (Iowa, United States). *Cancer Causes Control*; **9** (4), 357-367.
- Steinmetz KA, Kushi LH, Bostick RM, Folsom AR, Potter JD. (1994) Vegetables, Fruit, and Colon-Cancer in the Iowa Womens Health Study. *American Journal of Epidemiology* **139**(1), 1-15.
- Terry P, Hu FB, Hansen H, Wolk A. (2001) Prospective study of major dietary patterns and colorectal cancer risk in women. *Am J Epidemiol* **154** (12), 1143-1149.
- Wakai K, Date C, Fukui M, Tamakoshi K, Watanabe Y, Hayakawa N et al. (2007) Dietary fiber and risk of colorectal cancer in the Japan collaborative cohort study. *Cancer Epidemiol Biomarkers Prev* **16** (4), 668-675.
- Willett WC, Stampfer MJ, Colditz GA, Rosner BA, Speizer FE. (1990); Relation of meat, fat, and fiber intake to the risk of colon cancer in a prospective study among women. *N Engl J Med* **323** (24),1664-1672.

Annex 1: Carbohydrate definitions relating to fibre

Oligosaccharides- short chain carbohydrates comprising of 3-9 monomers joined together by glycosidic linkages.

Alpha glucans- oligosaccharides with an alpha linkage between monomers

Inulin, Fructooligosaccharides (FOS)- non- α - glucan oligosaccharides, known as fructans, and are the storage component of artichokes and chicory.

Polydextrose -non- α -glucan oligosaccharide

Galactooligosaccharides (GOS)- oligosaccharides found in milk, which principally contain galactose.

Non starch polysaccharide (NSP)- non- α -glucan polysaccharides that are mainly found in the plant cell walls. This includes cellulose, hemicellulose, pectin, arabinoxylans, plant gums, β -glucans.

Pectin- an NSP which is common to all cell walls.

Guar gum- an NSP which is which is chemically related to the cell wall, but is not strictly a cell wall component. Plant gums are sticky exudates which are formed at the site of injury.

Starch- the storage carbohydrate of plants, such as cereals, root vegetables and legumes, and consists of only glucose molecules.

Soluble fibre- relates to NSP components which can be rendered soluble by changing the pH conditions. These generally undergo significant fermentation, and viscous forms of these may also slow rates of glucose and lipid absorption from the small intestine^A. Examples of soluble fibres include pectin, beta-glucan (from oats and barley) and psyllium.

Insoluble fibre- NSP components that tend to undergo slow and incomplete fermentation and can have a greater effect on bowel habit^A. Insoluble fibres are found in vegetables and wholegrain products.

Resistant starch- the sum of starch and products of starch digestion (such as maltose, maltotriose and α -limit dextrins) that are not absorbed by the small bowel.

Prebiotic- a non-digestible food component that stimulates the growth and/ or activity of the bacteria in the bowel.

Lignin- non-carbohydrate component associated with plant walls

Englyst method- specifically determines NSP using an enzymatic-chemical method, which can be modified to yield soluble and insoluble fractions.

American Association of Analytical Chemists (AOAC)/ enzymatic-gravimetric method-

Determines total, soluble and insoluble residue containing carbohydrate and non-carbohydrate material in unknown proportions by measuring total residue weight and subtracting ash and protein content.

^AThe division between soluble and insoluble fibre is extremely pH dependent. Also, a large proportion of insoluble fibres are completely fermented and not all soluble fibre have effects on glucose and lipid absorption. Therefore, WHO have considered these definitions to be less useful when characterising fibre components.