



Paper for discussion : Update on Vitamin D

Agenda item: 6

This is an updated version of SACN/06/27, last seen by members at the October 2006 meeting. Members are asked to discuss and prepare a position statement.

This draft incorporates comments received from members. A summary table of comments and actions taken can be seen in Annex A.

Summary

- 1) Vitamin D is essential to the achievement of calcium homeostasis. It is found in a few foods, such as oily fish, fortified margarines and breakfast cereals and smaller amounts in red meat and egg yolk, but the main source in man is skin photosynthesis following irradiation with short wavelength ultra violet light (290-310 nm).
- 2) The National Diet and Nutrition Survey (NDNS) provides evidence of low vitamin D status in most age groups in the UK population, especially older children and young adults, and in older people living in institutions. Young women of childbearing age also have low vitamin D status and are likely to begin their pregnancies with low stores. Other evidence highlights a greater risk of vitamin D deficiency in population subgroups, particularly infants from black and ethnic minority groups. Cases of clinically apparent vitamin D deficiency in UK children, predominantly of Afro-Caribbean or South Asian origin, are widely reported but there is a lack of NDNS data for these population subgroups.
- 3) The Dietary Reference Values (DRVs) as defined in the 1991 COMA report do not set a Reference Nutrient Intake for vitamin D for adults or children over four years of age who receive adequate sunlight exposure. The current Reference Nutrient Intake for pregnant and breast-feeding women is 10µg vitamin D per day. For children under the age of four years it is 7-8.5µg per day and for those in the population aged over 65 years or confined indoors is 10µg vitamin D per day. These intakes will not be met from the diet and can only be guaranteed by medicinal supplementation. Although recommendations are available, there is concern that such recommendations are overlooked or not implemented by health professionals and the general public.
- 4) Deficiency of vitamin D results in the clinical illnesses of rickets (caused by osteomalacia) in children and osteomalacia in adults. The incidence of rickets in the UK declined from the 1920s onwards, which can partly be attributed to better living conditions, a reduction in atmospheric pollution, changes in diet, mandatory fortification of margarine with Vitamin D and replacement of cow's milk by infant formula during the first year of life. Rickets is now reported rarely among the white UK population although there is evidence of rickets in UK South Asian and Afro-Caribbean groups.. There is also recognition of a high prevalence of low vitamin D status among older people, particularly those living in institutions.
- 5) Several factors potentially affect vitamin D status. These include genetic factors, adiposity and factors affecting the cutaneous synthesis of vitamin D such as skin pigmentation, age, season, latitude, melanin concentration, clothing and use of sunscreens. Seasonal variations in vitamin D status are observed in the UK where the 2000/1 NDNS reported average plasma 25(OH)D concentrations to be highest in July-September and lowest in January-March. During winter, the UK population relies on body stores and dietary vitamin D to maintain vitamin D status. Solar UV radiation varies with latitude, and in winter months at latitudes of about 52° and above, there is no ultraviolet light of the appropriate wavelength for the cutaneous synthesis of vitamin D. For the remaining months, more than half the effective UV radiation occurs between certain times (1100 and 1500 hours) and is lower in the north than the south. Skin exposure to UV irradiation of

the appropriate wavelength is essential for maintaining adequate vitamin D status and a clear recommendation on length and intensity of exposure is required.

- 6) Although contradictory evidence exists concerning vitamin D status and subsequent implications for osteoporosis and fracture risk, a recent review of evidence suggests that a vitamin D intake of 17-20 μ g/d combined with calcium supplementation is associated with fewer hip and other non-vertebral fractures in frail older people confined to institutions. However, evidence for the effectiveness of vitamin D supplementation alone in fracture prevention remains unclear.
- 7) It is now recognized that the vitamin D receptor is expressed in most cells of the body and is involved in the regulation of gene transcription. There is accumulating evidence that vitamin D serves a modulatory role in a range of physiological processes in addition to calcium homeostasis. A low vitamin D status has been implicated in a range of other diseases including several forms of cancer, cardiovascular disease, tuberculosis, multiple sclerosis and type I diabetes. Research in these areas is developing, but evidence is inconclusive at present, and further work is needed before any definitive conclusions can be drawn.
- 8) Serum 25(OH)D concentration is used widely as a biochemical index of vitamin D status but there is no consensus on the threshold concentration that is diagnostic of a vitamin D deficiency state. Lack of functional criteria of sub-clinical deficiency and standardization of laboratory methodology makes risk assessment problematic. The development of standardized methodologies, together with a comprehensive review of the relationship between biochemical vitamin D status and health outcomes is required.
- 9) In view of these uncertainties, dietary supplements are necessary to prevent vitamin D deficiency in vulnerable subgroups of the population. The Committee reiterates the original COMA recommendation for vitamin D supplementation in vulnerable groups of the UK population including infants and young children, older adults, pregnant and breastfeeding women and black and ethnic minority groups, to safeguard vitamin D status.

Introduction

- 10) This paper highlights the re-emergence of rickets in population subgroups and the high prevalence of low vitamin D status throughout the UK population. The 1991 UK Dietary Reference Values (DRV) recommended that certain at-risk individuals or groups receive 7-10 μ g daily vitamin D. A Reference Nutrient Intake of 10 μ g/d was set for pregnant and lactating women, for the majority of whom supplementation will be required. Recent antenatal guidance from the National Institute for Health and Clinical Excellence, however, stated that there was insufficient evidence to evaluate the effectiveness of vitamin D supplementation in pregnancy and that, in the absence of evidence of benefit, vitamin D supplementation should not be offered routinely to pregnant women (National Institute for Health and Clinical Excellence, 2003). Discrepancy between these recommendations has led to a lack of clarity, which requires resolution to ensure clear guidance to health professionals and the general public.
- 11) This report is not a systematic review of Vitamin D and health but provides an update on vitamin D status and other related issues including a synopsis of evidence about the


relationship between vitamin D status and chronic disease, to assess the need for full consideration of the evidence and a full risk assessment.

Background

Vitamin D metabolism

- 12) Vitamin D is a pro-hormone that is produced photochemically in the skin from 7-dehydrocholesterol; however, if there is insufficient endogenous synthesis, generally caused by limited exposure of the skin to sunlight, then a dietary supply becomes essential. The action of sunlight (ultraviolet (UV) radiation of wavelength 290-310nm) on the skin converts 7-dehydrocholesterol to previtamin D₃, which is then metabolized to vitamin D₃ by isomerization. Vitamin D₃ is transported by vitamin D-binding protein to the liver. Dietary vitamin D exists as either ergocalciferol (vitamin D₂) or cholecalciferol (vitamin D₃), is fat-soluble and once ingested is incorporated into the chylomicron fraction, absorbed through the lymphatic system and transported to the liver. The liver enzyme 25-hydroxylase converts dietary and endogenously synthesized vitamin D to 25 hydroxyvitamin D (25(OH)D). Further conversion to the active form, 1,25-dihydroxyvitamin D (1,25 (OH)₂D), by 25(OH)D 1 α -hydroxylase (1 α -OHase) occurs under the influence of PTH in the kidney (DeLuca, 2004) (see Figure 1).
- 13) The only difference between ergocalciferol and cholecalciferol is the structure of the side chain to the sterol skeleton. Ergocalciferol is derived from the UV irradiation of the plant sterol ergosterol, which is widely distributed in plants and fungi; whereas, cholecalciferol is formed from the action of UV irradiation on skin. The widespread assumption that ergocalciferol and cholecalciferol are equipotent medicinally has recently been questioned (Houghton and Vieth, 2006).
- 14) The main circulating vitamin D metabolite is 25(OH)D and its concentration in plasma or serum is used as an indication of body status. The plasma concentration of 1,25(OH)₂D, the active form of the vitamin involved in calcium metabolism, is homeostatically regulated (Holick, 2004). The plasma 25(OH)D concentration is about a thousand-fold higher than 1,25(OH)₂D concentration, and provides a substrate reservoir for 1 α -OHase. The appearance in plasma of the parent compound, vitamin D, is short-lived since it is either taken up by adipose tissues or metabolized in the liver (Mawer *et al.*, 1972). The half-life of plasma 25(OH)D is about 2-3 weeks (Lund *et al.*, 1980); whereas, the half-life of plasma 1,25(OH)₂D is less than four hours (Holick, 2004).

Figure 1. The chemical conversion of vitamin D to the active hormone.



15) Vitamin D binding protein is a multi-functional plasma protein that transports vitamin D and its metabolites in the blood. It is a member of a gene family that includes albumin and alpha-fetoprotein and is identical to the group specific component (Gc-globulin) of serum. Vitamin D binding protein is synthesized in the liver and circulates at a concentration which is in excess of normal circulating vitamin D metabolite concentrations (Haddad, 1995). Like other fat-soluble compounds, only a small fraction of any vitamin D metabolite is freely dissolved in plasma. Vitamin D binding protein has a higher affinity for 25(OH)D than 1,25(OH)₂D (Bikle *et al.*, 1985; Bikle *et al.*, 1986; Teegarden *et al.*, 1991) and it is the free fraction of 1,25(OH)₂D that is functional *in vivo* (Vieth, 1990).

Vitamin D function

16) 1,25(OH)₂D acts in concert with parathyroid hormone and calcitonin to maintain serum calcium concentration within the normal range. This is achieved by regulating the efficiency of the small intestine to absorb calcium from the diet, by promoting the mobilization of calcium from the skeleton and by increasing the tubular reabsorption of calcium within the kidney (DeLuca, 2004). PTH and 1,25(OH)₂D together stimulate osteoblasts to induce the maturation of preosteoclasts to osteoclasts, thereby increasing bone resorption.

17) The synthesis of 1,25(OH)₂D in the kidney is tightly regulated, principally through the action of PTH. Calcium-sensing proteins in the parathyroid gland stimulate PTH secretion in response to a fall in serum calcium concentration. PTH promotes the renal synthesis of 1,25(OH)₂D₃, which, in turn regulates the synthesis of PTH by negative feedback (DeLuca, 2004).

18) An increase in serum 25(OH)D concentration, when serum 1,25(OH)₂D concentration remain unchanged, has been associated with suppression of serum PTH concentration and

an increased efficiency of calcium absorption (Heaney, 2004). It has been suggested that the local production of 1,25(OH)₂D (acting in an autocrine or paracrine fashion), as well as the endocrine function attributable to renal production, is important in many physiological processes (Fleet, 2004).

- 19) The active form of vitamin D, 1,25(OH)₂D, interacts with a specific receptor (VDR) similar to other steroid hormones. This receptor acts through nuclear vitamin D-responsive elements, which are usually found within 1 kilobase of the start site of the target gene and are involved in the regulation of gene transcription. VDR is expressed in cells involved in calcium homeostasis, e.g. enterocytes, osteoblasts, parathyroid and distal renal tubule cells, but is also expressed in cells unrelated to calcium homeostasis. VDR is present in the small intestine, colon, osteoblasts, activated T and B lymphocytes, pancreatic β-islet cells, and most organs in the body, including brain, heart, skin, gonads, prostate, breast, and mononuclear cells (DeLuca, 2004).
- 20) The 1α-OHase, which catalyses the conversion of 25(OH)D to 1,25(OH)₂D, has been shown to be expressed in many extrarenal tissues, including osteoclasts, skin, macrophages, placenta, colon, brain, prostate, endothelium and parathyroid glands (Zehnder *et al.*, 2001; Zehnder *et al.*, 2002; Segersten *et al.*, 2002; Schwartz *et al.*, 2004; van Driel *et al.*, 2006). Extrarenal production of 1,25(OH)₂D appears to play an important role in cell differentiation, proliferation and immune function. Vitamin D may, therefore, be involved in other physiological processes, independently of calcium metabolism. In contrast to renal 1α-OHase, extrarenal 1α-OHase is known to be unresponsive to stimulation by PTH and there is a lack of feedback inhibition by 1,25(OH)₂D and relatively low levels of 1,25(OH)₂D-directed catabolic 24-hydroxylase activity (Ren *et al.*, 2005; van Driel *et al.*, 2006).

Assessment of vitamin D status

- 21) Serum 25(OH)D concentration is used to assess vitamin D status. Plasma or serum concentration of 1,25(OH)₂D, and particularly free 1,25(OH)₂D, is a measure of vitamin D hormone activity, but, because of tight physiological regulation and levels that are much lower than that of its precursor metabolite, it does not reflect vitamin D nutritional status.
- 22) Several units of measurement are used to describe vitamin D intake and plasma 25(OH)D concentration. For vitamin D intake, 1 µg of dietary vitamin D is equivalent to 40 international units (IU). For plasma 25(OH)D concentration, 2.5 nmol/l is equivalent to 1ng/ml.
- 23) Different methods may be used to measure serum 25(OH)D concentration but comparisons are complicated by a lack of standardization. Different laboratories, using different methods, have been shown to yield different results from the same sample (Heaney, 2004; Binkley *et al.*, 2004). The international Vitamin D Quality Assessment Scheme (DEQAS) was established in 1989 to monitor the performance of 25(OH)D assays and now has over 100 registered participants in 18 countries, including the UK. For samples containing only 25(OH)D₃, DEQAS has found that most commercial 25(OH)D methods available in 2004 were capable of giving results close to the target value, but that the results were highly operator- and laboratory-dependent (Carter *et al.*,

- 2004). To complicate further the interpretation of results, some assay methods failed to detect 25(OH)D₂ with the same efficiency as 25(OH)D₃ (Carter *et al.*, 2004; Leventis *et al.*, 2005). New high performance liquid chromatography methodologies may offer a more robust and reliable measure (Lensmeyer *et al.*, 2006).
- 24) Serum 25(OH)D concentration in rickets and osteomalacia ranges from the undetectable to around 20 nmol/l (Department of Health, 1998). A serum 25(OH)D concentration of 25 nmol/l has been used as a conventional cut-off for defining the lower limit of adequacy of vitamin D status (Department of Health, 1998); however, this approach has been questioned and there is currently no consensus on the definition or biochemical criteria of a vitamin D deficiency state (Pal *et al.*, 2003; Bates *et al.*, 2003). This is reflected in the different thresholds adopted by studies cited later.
- 25) It has been suggested, for example, that vitamin D insufficiency or hypovitaminosis D, without clinical signs or symptoms, occurs at a serum 25(OH)D concentration of less than 40 nmol/l (Hanley & Davison, 2005), but there is no agreed definition. Based on associations between serum 25(OH)D concentration and serum PTH concentration, calcium absorption, bone turnover markers, and bone mineral density, others have argued that a serum 25(OH)D concentration of greater than 75 nmol/l is more appropriate to define vitamin D sufficiency or physiologically optimal concentrations (Hollis, 2005; Bischoff-Ferrari *et al.*, 2006).
- 26) The concept of establishing a reference range for serum 25(OH)D concentration based on a threshold at which serum PTH concentration starts to rise, is complicated by the large variation between individuals, the observation that this threshold varied between 30 and 78 nmol/l in several studies, and the fact that that no threshold could be identified in some studies (Department of Health 1998; Bates *et al.* 2003.) Dietary calcium intake influences serum PTH concentration, and serum PTH may also influence the turnover of Vitamin D metabolites (Lips, 2004). Inter-laboratory variation in determining serum 25(OH)D concentration may also influence the definition of a reference range.
- 27) The inverse relationship between serum 25(OH)D and PTH concentrations appears to be more pronounced with increasing age. Secondary hyperparathyroidism, associated with diminishing renal function, is observed in older people with poor vitamin D status (Vieth *et al.*, 2003). In a study of 1741 adults, those aged over 70 years had a higher mean PTH concentration than those aged less than 50 years and the plasma 25(OH)D concentration associated with a minimal PTH concentration was higher in the older subjects (Vieth *et al.*, 2003). The relationship between plasma 25(OH)D and PTH concentrations was investigated in the National Diet and Nutrition Survey (NDNS) of people aged 65 years and over (Bates *et al.*, 2003). An inverse association was observed between plasma 25(OH)D and PTH concentrations, but there was no evidence of a threshold for plasma a 25(OH)D concentration above which an irreducible minimum plasma PTH concentration was achieved.
- 28) Normative or optimal ranges for plasma 25(OH)D concentration have been suggested, based on associations with chronic disease end-points in older adults, e.g. osteoporosis and colorectal cancer (Dawson-Hughes *et al.*, 2005; Bischoff-Ferrari *et al.*, 2006). This is discussed further in the vitamin D and disease section.

Dietary sources of Vitamin D

- 29) There are few dietary sources of vitamin D. In the UK, rich sources are oily fish (5-10µg/100g), e.g. salmon, mackerel, sardines and fortified foods, e.g. margarines (~7µg/100g) and breakfast cereals (3-8µg/100g) (see Vitamin D food fortification section below). Red meat (~1µg/100g) and egg yolk (~5µg/100g) also provide vitamin D.
- 30) Supplemental vitamin D contains either ergocalciferol or cholecalciferol. Studies in the 1930s did not show any differences in antirachitic activity between the two forms, but more recent studies have shown that cholecalciferol appears to increase serum 25(OH)D concentrations more efficiently than does ergocalciferol (Tjellesen *et al.*, 1986; Trang *et al.*, 1998; Armas *et al.*, 2004). A lower binding affinity of ergocalciferol metabolites to the vitamin D binding protein in plasma may be a factor in these differences (Houghton & Vieth, 2006).
- 31) Human breast milk contains low concentrations of vitamin D and its metabolites. Exclusive breast-feeding for periods longer than 6 months in the absence of skin sunlight exposure of infants at the appropriate wavelengths or supplemental vitamin D, is associated with vitamin D deficiency, particularly if maternal stores were low throughout pregnancy (Ahmed *et al.*, 1995; Mughal *et al.*, 1999; Ziegler *et al.*, 2006).

Vitamin D supplements

- 32) Vitamin supplements provided to low-income families under the Welfare Food Scheme included 7 µg/day of vitamin D for children under 5 and 10 µg/day for pregnant and lactating women. However, the continued supply of these supplements has been problematic over the past few years. A reformed scheme for provision of welfare foods, Healthy Start, was launched in November, 2006; this provides for beneficiaries licensed supplements containing vitamins A, C and D for children and vitamins C, D and folic acid for pregnant and breastfeeding mothers.
- 33) Most commercial multivitamin preparations contain vitamin D but are deemed unsuitable for pregnant women because of their vitamin A content. No single vitamin D supplements are currently available in the recommended dose of 10µg/day, although this dose is available in supplements in combination with calcium.

Vitamin D food fortification

- 34) In the UK, manufacturers have practised voluntary fortification of margarine with vitamins since 1925. In 1940, with the advent of war, the Government took action to safeguard the nutritional status of the nation and the addition of vitamins A and D to all margarine sold for domestic use was made compulsory. This mandatory fortification was justified by evidence that a large proportion of the population, particularly children, were at risk of deficiency. Levels of added vitamin A were chosen to equate to the levels found in butter, but it was felt that vitamin D levels should be higher since margarine in the diet was considered to be the easiest, and possibly the only, means of ensuring an adequate supply of this vitamin to children in some sections of the community. Fat spreads, which have a fat content of less than 80%, are not subject to specific legislation, but some brands are fortified voluntarily to the same levels as margarine (Department of

Health, 1991b). This policy of fortification, together with other factors such as the later Clean Air Acts, improved housing, increasing affluence and longer and more frequent holidays, appeared to have been effective as rickets were very rare. Enquiries by the Ministry of Health between 1963 and 1966 then confirmed a low prevalence of rickets in some industrial cities, notably Glasgow (Department of Health, 1991b).

35) In 1952 a newly-recognized disease of infants and young children was described, with failure to thrive and occasionally fatal hypercalcaemia. Excessive intakes of vitamin D were suggested as a causative factor. At that time, cod liver oil compounds, infant milks and cereals were all fortified with vitamin D. The Ministry of Health and the Department of Health for Scotland concluded in 1957 that infants had unnecessarily high intakes of vitamin D and the levels in cod liver oil compound, infant milks and cereals, were reduced (British Paediatric Association, 1956; Ministry of Health and Department of Health for Scotland, 1957). In 1960, the incidence of idiopathic hypercalcaemia was believed by many British paediatricians to have decreased since reduction of the vitamin D content of various infant foods and supplements. Although it was generally inferred that a causal relation between vitamin D and infantile hypercalcaemia had been established, rigorous epidemiological evidence of a reduction in incidence was not available at that time (Fraser, 1967).

36) Early in the 1970s there were reports of vitamin D deficiency rickets and osteomalacia (see

37)

38) In a risk assessment by Hathcock *et al* (2007) based on well designed clinical trials of vitamin D in which doses in a range of 50 – 2500 µg/day were used, data was sufficient to show that vitamin D is not toxic at intakes much higher than previously considered unsafe. The authors concluded that in trials conducted in healthy adults that used vitamin D dose >250 µg/day, the absence of toxicity supports the selection of this dose as an upper limit (Hathcock *et al*, 2007).

39) Vitamin D deficiency, section below) in the UK, particularly in South Asian immigrants. COMA, therefore, convened a Working Party on Fortification of Food with Vitamin D which reported in 1980 (Department of Health and Social Security, 1980), which confirmed that rickets and osteomalacia did occur in South Asian children and women. The Working Party recommended that appropriate dietary supplements should be used and, while they did not recommend that fortification with vitamins A and D be extended to any other foods, they recommended that the mandatory fortification of margarine with vitamin A and D should continue.

40) In the US and a number of other countries, milk is fortified with vitamin D. In Finland in February 2003, the vitamin D fortification of liquid milk products (0.5 µg/dl milk) and margarines (10 µg/100 g) was introduced. A study of 196 young Finnish men (aged 18–28 years) observed that the prevalence of a serum 25(OH)D concentration <40 nmol/l decreased by 50%, from 78% in January 2003, before fortification to 35% in January 2004 (Laaksi *et al.*, 2006).

41) However, another study (Valimaki *et al.*, 2006) observed that vitamin D fortification in Finland improved the vitamin D status of 65 young Finnish men only marginally during winter: 33.8% of men had a serum 25-OHD concentration ≤ 20 nmol/l prefortification compared to 29.2% postfortification. However, the median serum 25-OHD concentration

prefortification was 24 nmol/l compared to 27 nmol/l postfortification.

- | [42](#)) Reports of hypervitaminosis D in the US caused by the inadvertent over-fortification of milk (Jacobus et al., 1992; Blank et al., 1995) have highlighted the need for careful monitoring of the effects of fortification programmes.

Exposure to sunshine

- | [43](#)) It is thought that most people in the UK obtain the majority of their vitamin D by exposure of skin to sunlight (Department of Health, 1998). The skin has a large capacity to produce vitamin D and exposure of about 20% of the body's surface to either direct sunlight or tanning bed radiation was effective in increasing the plasma concentration of 25(OH)D in both young adults and older adults (Holick, 2004).

- | [44](#)) The exact relationship between skin exposure to sunlight of the necessary wavelength and subcutaneous vitamin D synthesis is not well defined. In Cincinnati (latitude 38° N) during the spring, summer, and autumn, infants aged less than six months spending 20 minutes a day out of doors with exposed hand and face maintained 25OHD concentration above 27.5 nmol/l in infants aged less than 6 months (Specker *et al.*, 1985). It is suggested that exposure to sunlight for 5-15 min between the hours of 10:00 and 15:00 during the spring, summer, and autumn at latitudes above 37° is adequate for individuals with lighter skin (Holick, 2004).

- | [45](#)) The WHO/Euroskin Workshop on Vitamin D and UVR met in 2005, to discuss the growing controversy around how best to optimize vitamin D status while minimising the risk of diseases associated with sunshine exposure. The Workshop concluded that sun exposure is responsible for a substantial burden of skin and eye disease, and may play a role in reactivating some viral diseases. They also recognized that inadequate vitamin D status is a serious health issue but that more research is needed to establish optimal vitamin D levels for different groups within the general population. The solar UV index is an international standard measurement of how strong the ultraviolet radiation from the sun is at a particular place on a particular day. The group suggested that use of the solar UV index might be beneficial, where reference to location (latitude) and the likely UV sensitivity (skin type) of the recipient in articulating the need for varying levels of protection. It was agreed that the index should be actively promoted and form part of a sun protection programme, but presented in such a way as to enable individuals to relate the index to their own skin in interpreting what protective actions might be appropriate. There was general agreement that moderation of sun exposure is an important goal and that the key message is to ensure that people protect themselves when Solar UV Index is greater than 3 (McKinlay, 2006).

Factors affecting the cutaneous synthesis of vitamin D

Season and latitude

- | [46](#)) Solar UV radiation varies with latitude, time of year and time of day. From mid-October to the beginning of April at latitudes of about 52° and above (the UK is at latitude 50-60°N) there is no UV radiation of appropriate wavelength for the cutaneous production of previtamin D₃ (Webb & Holick, 1988). For the remaining months of the year 60% of the

effective UV radiation occurs between 11:00 and 15:00 hours, but is lower in the north than the south.

| [47](#)) Seasonal variations in plasma 25(OH)D concentration are observed in the UK. The 2000/1 National Diet and Nutrition Survey (NDNS) of adults aged 19 to 64 years reported average plasma 25(OH)D concentration to be highest in July-September (64.5 nmol/l for women; 64.9 nmol/l for men) and lowest in January to March (38.7 nmol/l for women; 40.8 nmol/l for men) (Ruston *et al.*, 2004). Summer plasma 25(OH)D concentration correlates with exposure to UV radiation and is largely a result of time spent outdoors as well as the amount of skin exposed (which can be subject to cultural influences) (Holick, 2004).

Formatted
Numbering

| [48](#)) During winter, the UK population relies on body stores and dietary vitamin D to maintain vitamin D status. The relative contribution of body stores and dietary intake is not well defined. In the UK, two studies of data from the NDNS of children and adults aged 65 years and over (Davies *et al.*, 1999; Bates *et al.*, 2003), observed the relationship between vitamin D intake and status to be seasonally dependent, being stronger in the winter and negligible in the summer. A study in healthy subjects in the US (Omaha; latitude 41.2° N) suggested that greater than 80% of the requirement for vitamin D in the winter months was met from body stores (Heaney *et al.*, 2003). A study in those aged over 65 years and over in the UK estimated that the plasma concentration during the summer months needed to be greater than 40 nmol/l to maintain plasma 25(OH)D concentration above 20 nmol/l during winter (Lawson *et al.*, 1979).

Formatted
Numbering

Age

| [49](#)) The amount of 7-dehydrocholesterol in the epidermis is relatively constant until later in life, when it begins to decline; a person 70 years of age exposed to the same amount of sunlight as a 20-year-old person makes about 25% of the vitamin D₃ that the younger person can make (Holick, 2004).

Formatted
Numbering

Skin pigmentation and ethnicity

| [50](#)) Skin pigmentation can also affect vitamin D₃ production, because melanin absorbs UV radiation in the 290-320nm range and functions as a light filter determining the amount of incident UV radiation available for the cutaneous production of previtamin D₃ (Norman, 1998). A study in Boston, USA (latitude: 42°N), observed mean plasma 25(OH)D concentration in young black women (n=51) to be less than half that in young white women (n=31) (Harris & Dawson-Hughes, 1998). Equally, whole body exposure to UV radiation resulted in a higher plasma 25(OH) D concentration in white and East Asian subjects than in South Asian or black subjects (Matsuoka *et al.*, 1991). Another study observed that a higher dose of UV radiation was required to increase plasma 25(OH)D concentration in South Asian subjects than white subjects, but the capacity to produce vitamin D was no different in Asian than in white skin (Lo *et al.*, 1986). It has also been suggested that altered vitamin D metabolism (increased 25(OH)D-24 hydroxylase activity) may be responsible for the high prevalence of low 25(OH)D concentration observed in South Asians (Awumey *et al.*, 1998).

Formatted
Numbering

Skin covering, and the avoidance of direct sun exposure

- 51)** Laboratory studies have shown that clothing prevents or significantly impairs the formation of vitamin D₃ after photostimulation (Matsuoka *et al.*, 1992). A study in Lebanon measured serum 25(OH)D concentration in 316 young adults (99 men and 217 women) from rural and urban areas. Almost three quarters of the study population had a concentration below 25 nmol/l, which was more common in women than in men, particularly in veiled women. In a multiple regression analysis; vitamin D intake, dwelling, veiling, and parity were independent predictors of hypovitaminosis D (Gannage-Yared *et al.*, 2000).
- 52)** A study in 321 healthy Saudi Arabian females showed that 52% had a 25(OH)D concentration ≤ 20 nmol/l (Ghannam *et al.*, 1999). This study did not specify the type of clothing, but a study in Kuwait showed that veiled women had lower 25(OH)D concentration than non-veiled women (el Sonbaty & Abdul-Ghaffar, 1996).

Sunscreen use

- 53)** Sunscreens are designed to prevent the entry of UV radiation before it enters the skin. Sunscreen with a sun protection factor (SPF) of 8 was shown to reduce the capacity of the skin to photoisomerize 7-dehydrocholesterol to previtamin D₃ by more than 95% in one laboratory based study (Matsuoka *et al.*, 1987) and, in a preliminary study, long-term users of sun screening agents were observed to have a lower serum 25(OH)D concentration than non-users (Matsuoka *et al.*, 1988).
- 54)** It has been suggested that the regular use of sunscreens might impair vitamin D status (Holick *et al.*, 1995; Ness *et al.*, 1999). A prospective study of 24 sunscreen users and 19 controls over 2 years, observed seasonal variation in serum 25(OH)D concentration in all subjects with slightly lower winter concentration in the sunscreen users (Farrerons *et al.*, 1998).
- 55)** A randomized controlled trial of the daily use of sunscreen (SPF 17) versus placebo cream over a summer period in Melbourne, Australia (latitude 37°S) was conducted in 113 people aged 40 years (Marks *et al.*, 1995). No differences by age, sex, and skin type were observed in the change in serum concentrations of 25(OH)D and 1,25(OH)₂D over the study period. The authors suggest that over an Australian summer sufficient skin exposure to sunlight was received, probably through both the sunscreen itself and the lack of total skin cover at all times, to allow a similar level of vitamin D production between people who are recommended to use sunscreens regularly and those who are not. No person using sunscreen developed serum vitamin D levels below the reference range over the period of the study.

Atmospheric pollution

- 56)** The first evidence of the importance of sunlight for human health began with the industrial revolution in northern Europe, where people congregated in cities and lived in dwellings that were built in close proximity to each other. The burning of coal and wood polluted the atmosphere reducing direct exposure to sunlight and, as a result, many children living in these industrialized cities developed rickets (Holick, 2004).

- | [57](#)) A study in India, compared serum 25(OH)D concentration in children 9-24 months old, from two regions of Delhi: one renowned for high levels of atmospheric pollution, the other less polluted (Agarwal *et al.*, 2002). Children living in areas of high atmospheric pollution had a mean serum 25(OH)D concentration of 31 nmol/l, less than half that of those living in the less polluted area.

Formatted
Numbering

Factors affecting vitamin D status

Vitamin D and adiposity

- | [58](#)) Vitamin D is fat-soluble and can be stored in the body fat. Adipose tissue was found to be the major site where vitamin D accumulated in human tissues after injection of radioactive vitamin D₃ (Mawer *et al.*, 1972). It is thought that excess vitamin D₃ produced during exposure to sunlight, and that is not broken down in the skin, is stored in body fat.

Formatted
Numbering

- | [59](#)) A lower serum 25(OH)D concentration and a higher serum PTH concentration were observed in obese subjects compared with non-obese subjects (Compston *et al.*, 1981; Hey *et al.*, 1982; Bell *et al.*, 1985; Liel *et al.*, 1988; Zamboni *et al.*, 1988; Buffington *et al.*, 1993; Hyldstrup *et al.*, 1993; Need *et al.*, 1993; Wortsman *et al.*, 2000; Parikh *et al.*, 2004; Snijder *et al.*, 2005; Hypponen & Power, 2006), indicating that the larger body fat compartments in the obese individuals may sequester vitamin D. In obese subjects compared with non-obese subjects, exposure to the same amount of UV radiation raised plasma 25(OH)D concentration by only 50% (Wortsman *et al.*, 2000). The content of the vitamin D₃ precursor 7-dehydrocholesterol in the skin was not significantly different between obese and non-obese subjects and the percentage conversion to previtamin D₃ and vitamin D₃ was similar in both groups. Obesity did not, therefore, affect the capacity of the skin to produce vitamin D₃, but may have altered the release of vitamin D₃ into the circulation, due to more being deposited in subcutaneous fat in obese subjects.

Formatted
Numbering

- | [60](#)) Total body fat, as determined by whole body dual energy x-ray absorptiometry scan (Arunabh *et al.*, 2003; Snijder *et al.*, 2005) and bioelectrical impedance analysis (Looker, 2005), has been shown to be inversely associated with plasma 25(OH)D concentration in healthy subjects. Amount of leg fat was also more strongly related to 25(OH)D concentration compared with abdominal fat, which might support the concept that endogenously produced vitamin D is deposited particularly in the subcutaneous fat depot (Snijder *et al.*, 2005).

Formatted
Numbering

- | [61](#)) Several small studies have reported morbidly obese individuals to have a higher 1,25(OH)₂D serum concentration than non-obese individuals (Hey *et al.*, 1982; Bell *et al.*, 1985; Liel *et al.*, 1988; Zamboni *et al.*, 1988). However, a more recent study in 302 healthy adults (152 obese) reported 1,25(OH)₂D serum concentration to be lower in obese subjects and to be inversely associated with percent total body fat, as determined by whole body dual energy x-ray absorptiometry scan (Parikh *et al.*, 2004). Methodological differences may be a factor in the apparent discrepancy with the earlier studies.

Formatted
Numbering

Genetics

- | [62](#)) Several polymorphisms in the *VDR* gene have been investigated with regard to vitamin D

Formatted
Numbering

status. A UK study of 143 healthy British South Asians, aged 31–65 years old, found serum 25(OH)D concentration did not vary with VDR genotype (*ApaI*, *BsmI*, *FokI* and *TaqI* polymorphisms) (Ogunkolade *et al.*, 2002); however, a study in Finnish women (n=93) observed that homozygotes for the VDR gene *BsmI*-polymorphism had a higher winter serum 25(OH)D concentration than other genotypes (Laaksonen *et al.*, 2002). In a study of 185 adolescent French girls, homozygotes for two polymorphisms in the promoter region of VDR had a lower serum 25(OH)D concentration than other genotypes (d'Alesio *et al.*, 2005). Phenotypes of the vitamin D-binding protein (Gc-globulin) were associated with plasma 1,25(OH)₂D and 25(OH)D concentrations in 595 Danish postmenopausal women, being highest in Gc1-1, intermediate in Gc1-2, and lowest in Gc2-2 phenotype (Lauridsen *et al.*, 2005).

- | **63)** In a study of 33 subjects, duodenal expression of the calcium channel/transporter gene *TRPV6* was strongly positively associated with plasma 1,25(OH)₂D concentration in men, but not in women, suggesting that *TRPV6* expression was vitamin D dependent in men (Walters *et al.*, 2006). An analysis of the *CDX2*-site polymorphism in the VDR promoter region, showed that in the *GG* genotype, but not the *AG* genotype, duodenal *TRPV6* expression was positively associated with plasma 1,25(OH)₂D₃ concentration (Walters *et al.*, 2004).

Formatted
Numbering

Exposure to sunshine and skin cancer

- | **64)** Most cases of skin cancer are caused by damage from UV rays in sunlight. As part of its SunSmart campaign, Cancer Research UK (CRUK) advises avoidance of the summer sun between 11am and 3pm by covering up with suitable clothing and by using a high-factor sunscreen when shade or clothing are not practical options. Fairer skinned people, those with a family history of skin cancer and people with many moles are at greater risk of skin cancer and need to take more care in the sun.

Formatted
Numbering

- | **65)** CRUK recognizes that a balance must be found to reduce further increases in skin cancer rates while also allowing enough sun exposure for optimal concentrations of vitamin D. A meeting to discuss the amount of sunlight needed for optimum health in European populations was convened in October 2005. CRUK, together with its advisory group of experts, is currently working on a collaborative position statement (CRUK, 2006).

Formatted
Numbering

Dietary Reference Values for the UK

- | **66)** The dietary reference values (DRV) for vitamin D set in 1991 were based on the dietary amount required to maintain plasma 25(OH)D concentration above 20 nmol/l in winter to prevent vitamin D deficiency (Department of Health, 1991a). A subsequent review examined the evidence on the relationship between bone health and vitamin D status and recommended no change to the DRV (Department of Health, 1998).

Formatted
Numbering

Table 1. Reference Nutrient Intakes (RNI) for vitamin D ($\mu\text{g}/\text{d}$) (Department of Health, 1998)

Age	Males	Females
0-6 months	8.5	8.5
7 months to 3 years	7	7
4 years to 65 years	-	-
65+ years	10	10
Pregnancy		10
Lactation, 0-4 months		10
Lactation, 4+ months		10

67) For 4 to 65 year olds it is assumed that the action of summer sunlight will provide adequate vitamin D status, except for specific at risk groups who are not exposed to sufficient sunlight, e.g. women whose clothing conceals them fully, those who are confined indoors. The RNI for these at risk groups is $10\mu\text{g}/\text{d}$. For the majority of people in this group, as well as the majority of pregnant and lactating women, people aged 65 years or more, infants and children aged up to 3 years, vitamin D supplementation will be needed to achieve the RNI (Department of Health, 1998).

68) It has been suggested that a vitamin D supplement of $10\mu\text{g}/\text{d}$ during lactation is insufficient to raise breast milk vitamin D and infant $25(\text{OH})\text{D}$ concentrations to levels that would have anti-rachitic activity and that a higher intake is required (Hollis & Wagner, 2004).

Safe upper intake levels for vitamin D

69) The cutaneous conversion of 7-dehydrocholesterol to previtamin D_3 , which spontaneously isomerizes to cholecalciferol, is regulated so that prolonged sunlight exposure does not lead to excess production; both precholecalciferol and cholecalciferol can be photolysed to inert compounds. High doses of oral vitamin D supplements, however, have been shown to have toxic effects (Vieth, 2006).

70) The toxic effects of vitamin D excess are primarily related to the role of the biologically active free form, $1,25(\text{OH})_2\text{D}$, in the regulation of plasma calcium (Pettifor *et al.*, 1995). Excessive production of $1,25(\text{OH})_2\text{D}$ or greatly increased plasma $25(\text{OH})\text{D}$ (which may displace $1,25(\text{OH})_2\text{D}$ from its binding protein and/or stimulate paracrine/autocrine $1,25(\text{OH})_2\text{D}$ production) can lead to an elevated plasma concentration of calcium (hypercalcaemia), due partly to over-stimulated intestinal absorption and partly to excessive calcium mobilization from bone. Hypercalcaemia could also lead to an increased calcium excretion into urine (hypercalciuria) (Vieth, 1990). Patients with sarcoidosis are abnormally sensitive to vitamin D, due to uncontrolled conversion of the vitamin to its active form in the granulomatous tissue. Although the condition is

uncommon, it would be a potential hazard if affected individuals were to take supplementary vitamin D (Expert Group on Vitamins and Minerals, 2003). Increased risk of hypercalcaemia has also been associated with combined treatment with vitamin D and thiazide and related diuretics (Mehta, 2006).

| **71)** In the UK, the Expert Group on Vitamins and Minerals report on safe upper levels for vitamins and minerals concluded that, for guidance purposes only, a level of 25µg/d supplementary vitamin D would not be expected to cause adverse effects in the general population when consumed regularly over a long period.

Formatted
Numbering

| **72)** The EU Scientific Committee on Food Opinion from 2002, published as part of the European Food Safety Authority tolerable upper intake levels for vitamins and minerals report (Scientific Committee on Food & Scientific Panel on Dietetic Products, 2006) could not establish a no observed adverse effect level (NOAEL) or a lowest observed adverse effect level (LOAEL), due to uncertainty in the data. A tolerable upper intake level (UL) was established at 25µg/d for infants and children aged 10 years or less and 50µg/d for children aged over 11 years and adults.

Formatted
Numbering

| **73)** The US Standing Committee on the Scientific Evaluation of Dietary Reference Intakes set a UL for infants aged up to 12 months of 25µg/d and for children aged 1 to 18 years and adults a UL of 50µg /d (Institute of Medicine, 1997).

Formatted
Numbering

| **74)** There is some controversy over the recommended safe upper limits in Europe and the USA. It has been argued that published cases of vitamin D toxicity with hypercalcaemia, for which the serum 25(OH)D concentration and vitamin D dose are known, all involved an intake of ≥1000 µg/d and that the UL of 50 µg/d, in the US, is too low by at least 5-fold (Vieth, 1999; Vieth, 2006). However, any of these studies involved no more than four weeks of supplementation, which may not be enough time for a steady state to be achieved (Scientific Committee on Food & Scientific Panel on Dietetic Products, 2006).

Formatted
Numbering

| **75)** In a risk assessment by Hathcock *et al* (2007) based on well designed clinical trials of vitamin D in which doses in a range of 50 – 2500 µg/day were used, data was sufficient to show that vitamin D is not toxic at intakes much higher than previously considered unsafe. The authors concluded that in trials conducted in healthy adults that used vitamin D dose ≥250 µg/day, the absence of toxicity supports the selection of this dose as an upper limit (Hathcock *et al*, 2007).

Formatted
Numbering

Vitamin D deficiency

| **76)** Vitamin D deficiency impairs the absorption of dietary calcium and phosphorus, which results in poor mineralization of the skeleton (Holick, 2006). Severe vitamin D deficiency generally presents as rickets (osteomalacia) in children and osteomalacia in adults. Rickets and osteomalacia are conditions characterized by pathological defects in growth plate and bone matrix mineralization. Histologically, the end result is an increased quantity of un mineralised bone matrix (osteoid).

Formatted
Numbering

| **77)** Rickets refers specifically to poor mineralization where the defect occurs in the growth plates at the end of the long bones. In children, failure of bone mineralization gives rise to bone deformities; bones are painful and linear growth is reduced. In adults, bone pain and

Formatted
Numbering

tenderness are the most prominent features of osteomalacia and proximal myopathy may also develop. Rickets can be precipitated by dietary calcium or phosphorus deficiency and calcipenic and hypophosphataemic rickets have been observed in children who are vitamin D sufficient (Thacher *et al.*, 1999) – the use of the term ‘rickets’ below refers to vitamin D deficiency rickets only.

| **78)** Rickets is the commonest presentation of vitamin D deficiency in children, but vitamin D deficiency may also present with hypocalcaemic symptoms – usually seizures, but occasionally more serious manifestations such as cardiomyopathy (Ladhani *et al.*, 2004).

| **79)** A plasma 25(OH)D concentration below 20 nmol/l is seen in children with nutritional rickets and concentrations below 10 nmol/l are observed in adults with osteomalacia (Department of Health, 1998).

| **80)** Evidence suggests there is an interaction between retinol and vitamin D with antagonistic effects and which has implications for bone health. Recent epidemiological evidence suggests that post-menopausal women with long-term high intakes of vitamin A have an increased risk of hip-bone fracture. These findings are supported by animal data, which have indicated that retinol has a direct effect on bone, possibly via an interaction with vitamin D, and an effect on parathyroid hormone and therefore calcium metabolism (Expert Group on Vitamins and Minerals, 2003). The vitamin D status of certain population subgroups, e.g. older people confined indoors and people from ethnic communities wearing enveloping clothing, may be poor and high intakes of retinol might be of greater concern in these populations (Scientific Advisory Committee on Nutrition, 2005).

Reports of clinically apparent vitamin D deficiency among children in the UK

| **81)** Prior to 1900, rickets affected about two-thirds of infants in the UK with the incidence of rickets peaking at the end of the 1800s. The incidence of rickets declined dramatically from the 1920s onwards, which may be partly attributed to better living conditions and changes in diet. In the period 1926 – 1942, studies in the UK indicated a prevalence of radiological signs of rickets among young children of 2% to 8%. A survey of 23 areas in the UK in 1943, showed on average, around 2% of infants had radiological signs of rickets (Department of Health and Social Security, 1980). Margarine was compulsorily fortified with vitamin D in the 1940s and cod liver oil, originally given as part of the Welfare Food Scheme (WFS), was substituted with vitamin supplements in 1975 (Department of Health, 2002). National Dried Milk was also fortified and manufacturers soon followed suit by adding vitamin A and D to infant milks, rusks and cereals. In the 1970s targeted campaigns of vitamin D supplementation, predominantly in Asian communities, helped further reduce the incidence of rickets in the UK (Dunnigan *et al.*, 1985).

| **82)** The sporadic incidence of rickets has continued; however, there is no information on the national prevalence of clinically apparent vitamin D deficiency in the UK. An unpublished review undertaken by Kings College, London in the 1990s (at the request of DH) confirmed that the problem of vitamin D deficiency and rickets, when occurring, still remained predominantly a problem in Asian populations. Where rickets was identified it was associated with strict vegetarian diets and exclusive breast feeding for periods longer

than 6-months.

| **83)** Over the past few years there have been several reports of clinically apparent vitamin D deficiency in UK children (infants and adolescents) (Pal & Shaw, 2001; Shaw & Pal, 2002; Ashraf & Mughal, 2002; Crocombe *et al.*, 2004; Odeka & Tan, 2005; Shenoy *et al.*, 2005; Callaghan *et al.*, 2006; Zipitis *et al.*, 2006). Most, though not all, of the cases that occur in the UK are seen in patients of Afro-Caribbean or South Asian origin. Although skin pigmentation is a factor, other factors, such as diet and low exposure to sunlight (either by staying indoors or covering the skin) are also important. An increase in clinically apparent vitamin D deficiency has been reported even in countries where sunlight is plentiful (Robinson *et al.*, 2006), underlining the importance of these lifestyle factors. Prolonged exclusive breast feeding also appears to be important, as breast milk contains little vitamin D (Ahmed *et al.*, 1995; Mughal *et al.*, 1999). It should be noted, however, that there is the possibility of a detection bias in reporting the link between ethnic group and South Asian children, as most of the reports come from areas with high South Asian populations; furthermore, national data are available for South Asian but not black children.

Formatted
Numbering

| **84)** A study in the UK, compared dietary intakes of 62 cases of rickets and osteomalacia and 113 normal women and children (Dunnigan *et al.*, 2005). An inverse association was observed between meat consumption and risk of rickets and osteomalacia; the relationship was curvilinear and relative risk did not fall further at meat intakes above 60 g daily. Daylight outdoor exposure was also inversely associated, and dietary fibre intake positively associated, with risk of rickets and osteomalacia.

Formatted
Numbering

| **85)** In a child health clinic in Manchester, six cases of florid rickets were described in infants of Asian ethnicity aged 10 to 28 months between 1995 and 1997 (Mughal *et al.*, 1999). The clinic subsequently reported that a further 8-10 non-white children with florid rickets were identified at their unit each year between 2001 and 2002 (Ashraf & Mughal, 2002). In a study of 124 ethnic minority children aged 6 to 36 months, two were identified with rickets, giving a prevalence of 1.6% (Ashraf & Mughal, 2002).

Formatted
Numbering

| **86)** A prospective survey conducted from May 2000 to April 2001 in the West Midlands reported 24 cases of clinically apparent vitamin D deficiency among children aged 0-4 years (Callaghan *et al.*, 2006); only one child was of white ethnic origin, the rest were either of South Asian or Afro-Caribbean origin. It was estimated from census data that the overall incidence was 7.5 per 100,000 children, with children of South Asian origin having an incidence of 38 per 100,000 and of Afro-Caribbean origin children having an incidence of 95 per 100,000. A separate survey of children under the age of 16 years presenting to three Birmingham hospitals between June 2001 and June 2003 identified 65 cases.

Formatted
Numbering

| **87)** A study in three London hospitals examined 65 children who presented with either rickets or hypocalcaemia and a plasma 25(OH D concentration below 25 nmol/l; 39 children were of Asian origin, 24 Afro-Caribbean, and two were Eastern European. Forty five percent (n=29) had hypocalcaemic symptoms, of whom 55% (n=17) had no radiological evidence of rickets; 48 children had radiological evidence of rickets, with or without other clinical signs (Ladhani *et al.*, 2004). Children who presented with hypocalcaemia were either under the age of 2 years or in adolescence. The authors speculated that during rapid

Formatted
Numbering

bone growth hypocalcaemia develops before rickets can ensue.

- | **88)** A study at Leicester hospital observed significant numbers of south Asian mothers having vitamin D deficiency at the end of pregnancy, and substantial numbers of children having infantile and adolescent rickets, some of whom have extremely severe bony deformities (Shenoy *et al.*, 2005). Increasing numbers of hypocalcaemic newborns, presenting predominantly with seizures, were also reported.
- | **89)** A survey conducted at the Burnley Health Care NHS Trust between 1994 and 2004 identified 14 cases of children presenting with clinically apparent (hypocalcaemia, rickets) vitamin D deficiency (Zipitis *et al.*, 2006). Thirteen of the 14 patients were of South Asian origin. From 1994 to 2001 there were 3 cases, but from 2002 to 2004 11 cases were reported, highlighting the rising incidence of clinically apparent vitamin D deficiency in the British Asian child community. The incidence of vitamin D deficiency was 1 in 117 for the Trust's Asian population compared to 1 in 923 in the general population. A cost-benefit analysis concluded that a campaign of vitamin D supplementation targeted to the South Asian population would be cost effective. The rise in incidence of rickets, especially in British Asian children, observed in the study was attributed to the lack of maternal vitamin D supplementation.
- | **90)** In the US, an assessment of women participating in the WIC programme (Special Supplemental Food Programme for Women, Infants and Children) suggested that factors that may have contributed to the increase in referrals of children with nutritional rickets included more African American women breast-feeding, fewer infants receiving vitamin D supplements and mothers and children exposed to less sunlight (Kreiter *et al.*, 2000)

Prevention of deficiency

- | **91)** Many countries have attempted to prevent such problems by ensuring that vulnerable groups receive dietary supplements of vitamin D during critical periods, such as pregnancy, lactation and infancy.
- | **92)** The UK RNI for vitamin D for all pregnant and breast feeding women is 10 µg of vitamin D daily, and for breast fed babies is 7-8.5 µg daily from the age of six. It is particularly important that pregnant women should have adequate vitamin D to build up their own stores and fetal stores to ensure adequate supply to infants during the first six months of life. Vitamin drops for children under 5 years of age (included in the WFS and Healthy Start) contain 7 µg of vitamin D and supplements for pregnant and nursing mothers contain a daily dose of 10µg.
- | **93)** There is concern that there may be a tendency for these recommendations to be overlooked by health professionals (Callaghan *et al.*, 2006; Department of Health, 1998), as well as by the general public. The review of the WFS identified that uptake of vitamin drops in the UK is very low even amongst those entitled to receive free supplies. The review concluded that the provision of free vitamin supplements offers a simple and potentially effective means of preventing adverse nutritional outcomes, particularly rickets. Rickets remains evident in the UK and it is likely that the prevalence would increase among high risk groups if the Scheme were withdrawn (Department of Health, 2002).

94) It has been questioned whether relying on vitamin D supplements given to infants or vitamin D supplementation of formula feeds is adequate to overcome the impact of maternal vitamin D deficiency (Shaw & Pal, 2002). This is also reinforced in a recent survey by Callaghan *et al* (2006), which highlighted that recommendations for vitamin D supplementation were being ignored and that 50% of those presenting with hypocalcaemic convulsions were formula fed (Callaghan *et al*, 2006). Shaw and Pal (2002) concluded that the definition of 25(OH)D should be standardized in the UK and recommended <20 nmol/l (8 ng/ml) during winter and <25 nmol/l (10 ng/ml) during summer. The authors proposed that all pregnant South Asian women be advised to take a vitamin D supplement in order to ensure that fetal stores are optimized; either 400 IU daily from the booking clinic appointment or 1000 units daily during the third trimester. Alternatively they recommended a single dose of ergocalciferol (100 000– 200 000 IU) during the sixth or seventh month may be preferred to achieve best efficacy and compliance. They also concluded that all Asian infants, whether formula or breastfed should receive 400 IU vitamin D daily, or alternatively an annual regimen may provide a higher compliance rate (Shaw and Pal, 2002).

Formatted
Numbering

95) There appears to be lack of awareness in high risk groups of the recommendations to take vitamin D supplements (Allgrove, 2004; Shenoy *et al.*, 2005; Callaghan *et al.*, 2006); furthermore, an audit in the Leicester area reported that while health professionals were aware of the issue there was no clear policy to resolve it (Iqbal *et al.*, 2001).

Formatted
Numbering

96) Antenatal guidance from the National Institute for Health and Clinical Excellence states that there is not enough evidence to evaluate the effectiveness of vitamin D in pregnancy and, that in the absence of evidence of benefit, vitamin D supplementation should not be offered routinely to pregnant women (National Institute for Health and Clinical Excellence, 2003). This resulted in uncertainty among health professionals with regard to supplementation during pregnancy. The Chief Medical Officer subsequently endorsed the COMA vitamin D recommendations for vulnerable groups including (Department of Health, 1998) pregnant and nursing mothers, young children and older people (Chief Medical Officer, 2005).

Formatted
Numbering

Dietary intakes and vitamin D status in the UK

97) The 1998 COMA report on Bone Health concluded that children aged 0-3 years, pregnant and lactating women and people aged 65 years or older, all of whom are vulnerable to vitamin D deficiency, had low dietary intakes even when provision from supplements was included. The contribution of supplements appears to be minimal except for infants and young children.

Formatted
Numbering

98) The DRV committee (Department of Health, 1998) assumed that most people aged 4-64 years of age will receive enough vitamin D from exposure of their skin to sunlight. Within this group, however, there may be individuals who are at risk of vitamin D deficiency and require dietary vitamin D to maintain adequate status. For these 'at risk' groups, e.g. where sunlight exposure is restricted, the RNI is 10 µg/d.

Formatted
Numbering

99) Infants and young children

- 100)** The NDNS of 1.5-4.5 year olds (Gregory *et al.*, 1995) found that mean intake of vitamin D from all sources was 1.9 µg/d (median 1.1 µg). Vitamin D intake from dietary supplements increased average intake by about half for all children. There were no significant differences between boys and girls or between age cohorts in intakes from food or from all sources. Children aged 1.5-2.5 obtained 17% of their mean vitamin D intake from "other milk and products", which included infant formula (fortified with vitamin D); children aged 2.5-4.5 obtained only 2-5% from these sources. Vitamin D fortified spreads (26%) and breakfast cereals (17%) were the other main food sources of vitamin D for all children.
- 101)** Mean plasma 25(OH)D concentration was 68.1 nmol/l and there was no apparent association with age or sex. There was seasonal variability in concentrations – highest among children assessed between July and September and lowest among children assessed between January to March. Plasma 25(OH)D concentration was below 25 nmol/l in 1% of children. The relation between vitamin D intake and status was also seasonally dependent, being strong in the winter and negligible in the summer (Davies *et al.*, 1999).
- 102)** A population survey in 1996 of children aged 2 years (n=618) of Bangladeshi, Indian, or Pakistani origin living in England showed that 20–34% had a plasma 25(OH)D concentration (measured during October-November 1996) less than 25 nmol/l; 3-18% had a value below 20 nmol/l (Lawson & Thomas, 1999). A significant association was found between failure to take a vitamin supplement, chapatti consumption and low plasma 25(OH)D concentration (Lawson *et al.*, 1999).
- 103)** The 1998 COMA report on Bone Health concluded that “vitamin D status, assessed from plasma 25(OH)D, of the majority of the population of children under 4 years appears to be satisfactory. Some minority groups of children remain at risk due to factors associated with lifestyle. The current programme of vitamin D supplementation for this section of the population should continue. Education programmes to reinforce this policy appear to be needed.

Older children

- 104)** The NDNS of 4-18 year olds (Gregory *et al.*, 2000) showed that mean daily intake of vitamin D from food sources for boys was 2.6 µg, significantly higher than that for girls at 2.1 µg (medians 2.4 and 1.9 µg/d, respectively). Mean intake increased with age. The main dietary sources of vitamin D were cereal and cereal products (37% boys and 35% girls), fat spreads (20% both sexes), meat and meat products (20% both sexes) and oily fish (7% boys and 9% girls). Supplements of vitamin D increased mean intake from food sources by 8% for boys and 5% for girls. The younger children predominantly took supplements. In the 4 to 6 age group, supplements increased mean intake by 19% for boys and 22% for girls.
- 105)** Correlation between plasma 25(OH)D concentration and dietary intake of vitamin D was very weak and did not reach the level of statistical significance at any age or sex group. The mean plasma 25(OH)D concentration was 62.0 nmol/l for boys and 60.6 nmol/l for girls. Three percent of boys and 2% of girls aged 11-18 years had a plasma 25(OH)D concentration less than 12 nmol/l. The proportion below 25 nmol/l increased with age for both boys and girls from 3% for boys and 2% for girls aged 4 to 6 years to

10-16% for boys aged 11 to 18 and 10-11% for girls aged 11-18. Plasma concentration was influenced by season of sample collection, as for younger children. Boys aged 11 to 18 years were the most likely to have poor vitamin D status. From January to March there were 19% of boys in the age group with poor status, 15% from April to June, 6% from July to September and 10% from October to December.

106) A study in Manchester showed that of 51 adolescent girls (aged 14.7 to 16.6 years; 14 white; 37 non-white) 73% (n=37) had a serum 25(OH)D concentration below 30 nmol/l and 17% (n=9) below 12.5 nmol/l (Das *et al.*, 2006). Non-white girls (21 South Asian; 5 Middle Eastern; 1 black) were more severely vitamin D deficient. Reduced sunlight exposure, rather than diet, accounted for the differences in vitamin D status.

Formatted
Numbering

Pregnant and lactating women

107) There is evidence to link vitamin D deficiency in infancy to poor maternal vitamin D status. Vitamin D deficiency in pregnant mothers is associated with congenital rickets and craniotabes in the newborn (Specker, 1994) and with rickets in infancy, especially when the child is exclusively breast-fed (Mughal *et al.*, 1999; Kreiter *et al.*, 2000). A poor maternal vitamin D status at a level above that associated with rickets in the child can adversely affect fetal and infant skeletal growth and ossification, tooth enamel formation and calcium handling (Specker, 1994). The vitamin D status of the infant appears to be more influenced by the vitamin D status of the mother during pregnancy, and by the infant's sunshine exposure, than by maternal vitamin D status during lactation (Specker, 1994).

Formatted
Numbering

108) Human milk contains relatively small amounts of vitamin D, between 0.3 and 1.5 µg/l, and concentrations of vitamin D and 25(OH)D in breast milk correlate with those in the mother's circulation (but at lower concentrations). Although these correlations might predict that the mother's vitamin D status could be important in determining the infant's vitamin D status, this does not appear to be the case. Infant plasma 25(OH)D concentration does not correlate with milk 25(OH)D concentration (38), unless the mother receives high doses of supplemental vitamin D (Specker, 1994). A doses of 50 µg/d given to lactating mothers increased infant serum 25(OH)D concentration, yet a dose of 25 µg/d did not do so (Ala-Houhala, 1985; Ala-Houhala *et al.*, 1986); supplementation of infants with 10 µg/d vitamin D raised serum 25(OH)D concentration to a similar extent to maternal supplementation with 50 µg/d. In breastfeeding women a dose of 50 µg/d, and more so 100 µg/d, of vitamin D₂ and vitamin D₃ was shown to raise breast milk concentrations of vitamin D and 25(OH)D and infant serum 25(OH)D concentration (Hollis & Wagner, 2004). Maternal exposure to UV radiation of appropriate wavelengths has been shown to increase the anti-rachitic activity of human milk (Greer *et al.*, 1984) and, under normal circumstances, the sunshine exposure of breast-fed infants is the major factor affecting their vitamin D status (Specker, 1994).

Formatted
Numbering

109) The 1998 COMA Report on Bone health noted that there was no national information on the vitamin D status of pregnant and lactating women. A study of 160 pregnant women from non European ethnic minorities living in South Wales found that 50% of women had a plasma 25(OH)D concentration below 20 nmol/l on their first antenatal visit (Datta *et al.*, 2002). The study found that fluency in English, dressing habits and religion did not appear to influence status, but a higher proportion of women who had lived in

Formatted
Numbering

Britain for longer than three years had a low 25OHD concentration. A relatively high number of South Asian mothers in Leicester have also been reported to have vitamin D deficiency at the end of pregnancy (Shenoy *et al.*, 2005), although the authors do not indicate how this was assessed.

110) A study of white women and their offspring (n=198) in Southampton (latitude 50° N) (Javaid *et al.*, 2006), observed 31% of mothers had a serum 25(OH)D concentration between 27.5-50 nmol/l and 18% had a serum 25(OH)D concentration less than 27.5 nmol/l during late pregnancy. Mothers who had a plasma 25(OH)D concentration of less than 27.5 nmol/l in pregnancy had an offspring whose whole-body bone mineral content at 9 year's of age was lower than those born to mothers with a higher plasma 25(OH)D concentration. Both the estimated exposure to UVB radiation during late pregnancy and the maternal use of vitamin D supplements predicted maternal 25(OH)D concentration and childhood bone mass. A low concentration of umbilical-venous calcium also predicted lower childhood bone mass.

Adults aged 19-64 years

111) Mean daily intake of vitamin D from food sources was 3.7µg for men and 2.8µg for women. Inclusion of supplements containing vitamin D increased mean intake by 14% in men, to 4.2µg, and by 32% for women, to 3.7µg. For women aged 50-64 years supplements increased mean daily vitamin D intake by 46%, from 3.5µg to 5.1µg. Median daily intake from all sources, was 2.1µg for women (aged 19-64 years) living in a household receiving certain benefits, compared with 2.9µg for women not receiving benefits (Henderson *et al.*, 2003).

112) The NDNS of 19-64 year olds (Ruston *et al.*, 2004) showed that the mean concentration of plasma 25(OH)D was 48.3 nmol/l for men and 49.6 nmol/l for women. A plasma 25(OH)D concentration below 25 nmol/l was found in around 15% of the adult population overall and a quarter of the 19-24 age group. This proportion was higher during the winter months.

113) A study in Birmingham of 240 adults compared plasma 25(OH)D concentration in South Asians with non-Asians (white and Afro-Caribbean) (Pal *et al.*, 2003). South Asians had a lower plasma 25(OH)D concentration than non-Asians in both summer and winter; the majority of South Asians had a concentration below 30 nmol/l (94% in winter; 82% in summer).

Adults aged over 65 years

114) Mean vitamin D intake was below the RNI (10µg) in all groups: mean daily intake from all sources was 3.47µg for institutionalized and 3.92µg for free-living subjects. Overall, 6% of men and 10% of women in the free-living group had a plasma 25(OH)D concentration below 25 nmol/l; this increased in the winter months (Finch *et al.*, 1998).

115) Vitamin D status was significantly worse in the institution group than in the free-living group. Over a third of men and women had a plasma 25(OH)D concentration below 25 nmol/l and there was no evidence of seasonal variation. Vitamin D intake from food sources was below the RNI (10 µg/day) with mean daily intakes of 3.42µg for

institutionalized and 3.40µg for free-living subjects, and the contribution from supplements was low (Finch *et al.*, 1998). The relationship between plasma 25(OH)D concentration and vitamin D intake in free-living subjects was seasonal: 25(OH)D concentration was associated with vitamin D intake in the winter, but not in the summer (Bates *et al.*, 2003).

| 116 In the Health Survey for England 2000, serum 25(OH)D concentration was measured in 1,766 people aged 65 years and over living in private households and institutions (Hirani & Primatesta, 2005). The mean serum 25(OH)D concentration was lower for both men (38.1 nmol/l) and women (36.7 nmol/l) in institutions than among men (56.2 nmol/l) and women (48.4 nmol/l) in private households. In institutions, the prevalence of serum 25(OH)D concentration below 25 nmol/l was 30.2% for men and 32.5% for women, while in private households women (15.0%) had a higher prevalence of low 25OHD concentration than men (9.6%).

Formatted
Numbering

Summary

| 117 The NDNS provides evidence of low vitamin D status in most population age groups, especially older children and young adults, and in older people living in institutions. Figure 1 below shows the percentage of people from different age groups with a plasma concentration of 25-hydroxy D below 25 nmol/l. Almost a third of young women of childbearing age (19- 24 years) appear to have low status and are likely to start pregnancy with low maternal stores. Routine supplementation of this group of women will be necessary, as recommended by COMA, in order to ensure adequate foetal supply and stores in the newborn.

Formatted
Numbering

| 118 Other studies also highlight population subgroups, e.g. those of South Asian origin, that have an even higher high prevalence of low vitamin D status and in whom rickets is increasingly common.

Formatted
Numbering

Fig 2. Percentage in NDNS with a plasma 25(OH)D concentration <25 nmol /l. F, free living; I, institutionalized.





Fig 3. Percentage in NDNS with a plasma 25(OH)D concentration <40 nmol /l. F, free living; I, institutionalized. Data for 1.5-4.5 year olds not available.



Vitamin D and disease

Osteoporosis and fracture risk

119) Two meta-analyses of primary prevention supplementation trials (5 trials for hip fracture risk (n=9,294), 7 trials for non-vertebral fracture risk (n=9,820), and 5 trials for falls (n=1,237)) concluded that vitamin D supplements may have a beneficial effect on bone mineral density, fracture risk (Bischoff-Ferrari *et al.*, 2005) and falls (Bischoff-Ferrari *et al.*, 2004). The beneficial effects were observed in trials of older adults where the dose of vitamin D was between 17.5 and 20 μ g/d, but not at 10 μ g/d, and when serum 25(OH)D concentration was about 75 nmol/l or more.

120) Several trials have been published since these meta-analyses were conducted. A secondary prevention trial in people aged 70 years (n=5,292) or more who had had a low-trauma, osteoporotic fracture in the previous 10 years, found no evidence of a beneficial effect of 20 μ g/d vitamin D₃ over 2 years on fracture prevention (Grant *et al.*, 2005). Based on the observed rise in serum 25(OH)D concentration in a subset of subjects, and the recorded compliance rate, it has been suggested that a lack of subject compliance and low baseline concentrations resulted in serum 25(OH)D concentration being too low for an effect on fracture rate (Bischoff-Ferrari *et al.*, 2006).

121) In another trial conducted in the UK (Porthouse *et al.*, 2005) 3,314 women, aged 70 years and over with one or more risk factors for hip fracture, received either daily supplements of 1000 mg of elemental calcium with 20 μ g of vitamin D₃ and an information leaflet on dietary calcium intake and prevention of falls, or leaflet only for 25 months. No effect of supplementation on fracture rates was observed; although, the odds ratio for hip fracture was 0.75 the 95% confidence intervals were large (0.31 to 1.78). Serum 25(OH)D concentration was not measured in this trial. It has been suggested that

the open design of the trial and instruction to the control group to ensure adequate calcium and vitamin D intakes may have biased the result towards the null hypothesis (Bischoff-Ferrari *et al.*, 2006).

122) In a US trial among healthy postmenopausal women (n=36,282) involving supplementation with calcium and vitamin D (1000 mg/d of elemental calcium and 10µg/d of vitamin D₃) did not significantly reduce hip fracture rate, although hip bone mineral density was slightly improved. However, a detrimental effect of supplementation (increased risk of kidney stones) was observed in this study (Jackson *et al.*, 2006). Again, based on the observed rise in serum 25(OH)D concentration in a subset of subjects, it has been suggested that a higher dose of vitamin D₃ would have been required to see an effect on fracture rates (Bischoff-Ferrari *et al.*, 2006).

123) A recent review of the effects of vitamin D or analogues in the prevention of fractures in older people (Avenell *et al.*, 2006), indicated that frail older people confined to institutions may sustain fewer hip and other non-vertebral fractures if given vitamin D (17-20µg/d) with calcium supplements. However, there was no evidence of effect of vitamin D with calcium on vertebral fractures and effectiveness of vitamin D alone in fracture prevention was unclear.

Colorectal cancer

124) Vitamin D insufficiency has been implicated in the development of cancer (Garland *et al.*, 2006), especially colorectal cancer (Gorham *et al.*, 2005; Giovannucci, 2006). The evidence from prospective cohort studies that have investigated dietary vitamin D intake and colorectal cancer risk are mixed. Several studies found an inverse association (Garland *et al.*, 1985; Martinez *et al.*, 1996; McCullough *et al.*, 2003), whilst others either showed no association (Jarvinen *et al.*, 2001; Lin *et al.*, 2005) or an inverse association which was not significant after adjustment for confounding variables (Bostick *et al.*, 1993; Kearney *et al.*, 1996).

125) A small nested case-control prospective study (34 cases) in the US (Garland *et al.*, 1989), observed a serum 25-OHD concentration of 50 nmol/l or more to be associated with a reduced risk of colon cancer.

126) In a prospective nested case-control study in the US (Feskanich *et al.*, 2004), higher plasma concentration of 25(OH)D was associated with a lower risk of colorectal cancer in women aged 46 to 78 years (193 colorectal cancer cases), particularly for cancers at the distal colon and rectum. Several case-control studies determining the risk of colorectal adenoma in relation to serum 25(OH)D concentration have also reported an inverse association and based on the available epidemiological evidence it has been argued that a serum 25(OH)D concentration of ≥90 nmol/l would be optimal (Bischoff-Ferrari *et al.*, 2006).

127) A prospective study in the US (Giovannucci *et al.*, 2006) considered multiple determinants of vitamin D exposure (dietary and supplementary vitamin D, skin pigmentation, adiposity, geographic residence and leisure-time physical activity – to estimate sunlight exposure) in relation to cancer risk. Based on measurements in 1,095

men of this cohort, Giovannucci *et al* quantified the relation of these six predictors to plasma 25(OH)D level and used results from the multiple linear regression model to predict 25(OH)D levels for each of 47,000 men and prospectively examined this variable in relation to cancer risk. A predicted low plasma 25(OH)D concentration was associated with increased cancer incidence and mortality, particularly for digestive-system cancers. This study used an indirect assessment of plasma 25(OH)D concentration that could be influenced by established surrogates such as race and geographic residence.

| **128)** A trial among healthy postmenopausal women (n=36,282), calcium with vitamin D supplementation (1000 mg/d of elemental calcium and 10µg/d of vitamin D₃) for 7 years did not reduce the risk of colorectal cancer (Wactawski-Wende *et al.*, 2006). It has been suggested, however, that, based on the epidemiological evidence, the supplementation period was not long enough, nor the dose of vitamin D sufficient, for an effect to become apparent (Bischoff-Ferrari *et al.*, 2006).

Other cancers

| **129)** Some epidemiological evidence also suggests associations between low vitamin D status and risk of prostate and breast cancers, but the evidence is weak (Giovannucci, 2005). A review of case-control and cohort studies (Cui & Rohan, 2006) found no association between dietary vitamin D intake and breast cancer risk, but plasma 25(OH)D concentration was inversely associated with breast cancer risk in a case-control study (Lowe *et al.*, 2005), but not significantly in a prospective nested case-control study (Bertone-Johnson *et al.*, 2005).

| **130)** Pre-diagnostic serum 25(OH)D concentration has been assessed in relation to prostate cancer risk in several prospective nested-case control studies. A 3-fold increased risk was observed in Finnish men aged less than 50 years with serum 25(OH)D concentration of <40 nmol/l (Ahonen *et al.*, 2000). A study in Swedish and Norwegian men observed both high (≥ 80 nmol/l) and low (≤ 19 nmol/l) serum 25(OH)D concentrations to be associated with a higher prostate cancer risk (Tuohimaa *et al.*, 2004). Several prospective nested case-control studies conducted in the US have, however, observed no association between serum 25(OH)D concentration and prostate cancer risk (Corder *et al.*, 1993; Braun *et al.*, 1995; Gann *et al.*, 1996; Nomura *et al.*, 1998; Platz *et al.*, 2004). It has been suggested that the subjects in the US studies may have a higher mean vitamin D status than the subjects in the Nordic studies; subsequently, the increased risk associated with a low 25(OH)D concentration in the Nordic studies may be difficult to detect in the US studies (John *et al.*, 2005).

| **131)** A prospective study in the US, in cohorts of 46,771 men ages 40 to 75 years and 75,427 women ages 38 to 65 years, examined the relation between dietary vitamin D intake and pancreatic cancer risk (Skinner *et al.*, 2006). Compared with participants in the lowest category of total vitamin D intake (<3.75µg/d), pooled adjusted relative risks for pancreatic cancer were 0.78 (95% CI, 0.59-1.01) for 3.75-7.47/d, 0.57 (95% CI, 0.40-0.83) for 7.50 to 11.22 µg/d, 0.56 (95% CI, 0.36-0.87) for 11.25-14.97 µg/d/d, and 0.59 (95% CI, 0.40-0.88) for ≥15µg/d (P_{trend} = 0.01).

| **132)** In a prospective nested case-control study in male Finnish smokers, those in the highest quintile of serum 25(OH)D concentration (>65.5nmol/l) had a 3-fold increased

risk for pancreatic cancer relative to those in the lowest quintile (<32.0 nmol/l) (Stolzenberg-Solomon *et al.*, 2006).

Cardiovascular disease risk factors

133) Low vitamin D status has also been implicated in cardiovascular disease risk (Zittermann *et al.*, 2005), although the evidence is relatively weak compared to bone health and colorectal cancer (Bischoff-Ferrari *et al.*, 2006).

134) A trial, conducted in men with congestive heart failure who were randomly assigned to receive a placebo or vitamin D (50 µg/d), found no effect of vitamin D on either left ventricular function or 15-mo survival rates (Schleithoff *et al.*, 2006); however, the serum concentration of tumour necrosis factor-α, an inflammatory cytokine, decreased with vitamin D treatment; in contrast, the concentration of interleukin 10, an anti-inflammatory cytokine, increased.

135) One trial observed a beneficial effect of vitamin D₃ supplementation (33µg/d for one month) on insulin sensitivity in type 2 diabetics (Borissova *et al.*, 2003) and a study, in normoglycemic subjects, observed insulin sensitivity to be positively correlated with 25(OH)D concentration. A concentration below 50nmol/l was associated with reduced pancreatic β cell function (Chiu *et al.*, 2004). An association between type II diabetes and a low serum 25(OH)D concentration in British South Asians has also been observed (Boucher, 2006).

136) Two trials have reported a lowering effect of the active form of vitamin D on blood pressure in normocalcaemic (Lind *et al.*, 1988b) and hypercalcaemic individuals (Lind *et al.*, 1988a). A trial giving either a single oral dose of 2.5 mg vitamin D₃ or placebo to 189 elderly individuals in the winter observed no effect on blood pressure nor serum cholesterol concentrations after 5 weeks (Scragg *et al.*, 1995). A trial in 148 elderly women, with a serum 25(OH)D concentration below 50 nmol/l, who received either supplemental calcium (1200 mg/d) or supplemental calcium (same dose) and vitamin D (20µg/d) for 8 weeks, observed that the group receiving calcium and vitamin D had a 9.3% decrease in systolic blood pressure relative to those receiving calcium alone (Pfeifer *et al.*, 2001). A large prospective study on over 200,000 subjects in the US, however, found no association between vitamin D intakes and hypertension (Forman *et al.*, 2005).

137) In 146 healthy, non-diabetic British South Asians both soluble C-reactive protein concentration and plasma metalloproteinase 9 concentration were inversely related to vitamin D status (Timms *et al.*, 2002). A higher plasma concentration of C-reactive protein metalloproteinase 9 has been associated with an increased risk of cardiovascular disease.

Tuberculosis

138) Vitamin D deficiency has been associated with susceptibility to tuberculosis (Ustianowski *et al.*, 2005), but paradoxically, tuberculosis has been associated with increased production of the active vitamin D metabolite, 1,25(OH)₂D₃ (Barnes *et al.*, 1989). TB-activated monocytes and macrophages account for increased 1,25(OH)₂D₃ production in and the increased vulnerability of subjects with vitamin D deficiency to

tuberculosis was explained by the requirement for sufficient 25(OH)D as substrate for 1,25(OH)₂D₃ production by the activated monocytes/macrophages (Liu *et al.*, 2006). Activation of Toll-like receptors (TLRs) during infection triggers direct antimicrobial activity against intracellular bacteria which up-regulates expression of the VDR and 1 α -OHase, leading to induction of the antimicrobial peptide cathelicidin and subsequent killing of intracellular *Mycobacterium tuberculosis*.

Other diseases

139) Prospective studies in Finland have observed a reduced risk of type I diabetes (Hypponen *et al.*, 2001) and an increased risk of atopy and allergic rhinitis (Hypponen *et al.*, 2004) in infants who regularly received vitamin D supplements during the first year of life relative to those who did not. A multi-centre study in the EU, also observed that vitamin D supplementation in infancy was associated with a decreased risk of type I diabetes (EURODIAB Substudy 2 Study Group, 1999).

140) A prospective study in the US (Munger *et al.*, 2004), observed vitamin D intake from supplements, but not foods, to be inversely associated with risk of multiple sclerosis. A subsequent prospective, nested case-control study observed the risk of multiple sclerosis to be inversely associated with serum 25(OH)D concentration (Munger *et al.*, 2006). Those in the lowest quintile of serum 25(OH)D concentration (<63.3 nmol/l) had a higher associated risk of multiple sclerosis than those in the highest quintile (>99.1 nmol/l).

141) A prospective study in the US (McAlindon *et al.*, 1996), reported low intake and low serum concentration of 25OHD to be associated with an increased risk for progression of osteoarthritis of the knee. Another prospective study in the US, reported that subjects with serum 25(OH)D concentration in the highest tertile had a reduced risk of incident changes of radiographic hip osteoarthritis, but not with incident hip osteoarthritis defined as the development of definite osteophytes or new disease (Lane *et al.*, 1999).

Conclusions

142) A significant proportion of the UK population have low vitamin D status based on the current COMA definition (a plasma 25-hydroxy vitamin D concentration below 25nmol/L). This increases their risk of vitamin D deficiency. This is a particular concern for women of child-bearing age, pregnant women, infants, the elderly and black and ethnic minority groups.

143) A lack of national data, for certain population subgroups, particularly South Asian and Afro-Caribbean; pregnant and breastfeeding women; infants, makes it difficult to estimate precisely the prevalence of low vitamin D status in the UK population. The Committee suggests the need for further surveys of vitamin D status in order to fully quantify the problem in the UK and to monitor prevalence into the future.

144) There is accumulating evidence on the association between vitamin D status and outcomes other than rickets and osteomalacia, indicating that the risks from low vitamin D status in the UK population may be greater than previously thought.

145) Recent evidence on the relationship between plasma 25OH vitamin D concentration

and other health outcomes suggests that the COMA threshold level of 25 nmol /L may be conservative. Further risk assessment and consideration of existing Dietary Reference Values is therefore warranted when more definitive evidence is available in the next 3-4 years from ongoing research by the Food Standards Agency.

| 146) In order to enable a full risk assessment of the impact of low vitamin D status, there is an urgent need to standardize laboratory methodologies for the measurement of serum 25(OH) vitamin D concentration. This will allow more robust assessment of the relationships between plasma 25(OH) vitamin D concentration and health outcomes in order to define threshold concentrations indicative of adequate population status.

Formatted
Numbering

| 147) In the meantime the current Dietary Reference Values for vitamin D set by COMA for pregnant and breast-feeding women, young children, people aged 65 years and over, and individuals who are at risk of inadequate sunshine exposure are endorsed. The committee reiterates COMA's recommendation that all pregnant and breastfeeding women should consider taking a daily supplement of 10µg vitamin D in order to ensure their own requirement for vitamin D is met and to build adequate fetal stores for early infancy.

Formatted
Numbering

| 148) The review of the Welfare Food Scheme revealed a tendency to overlook advice on vitamin D supplementation as there is a lack of awareness in vulnerable groups. The Committee also recognizes that there is generally a poor understanding of advice among health professionals and a lack of clarity in prescribing policy and the provision of appropriate supplements. A clear public health strategy and guidance on vitamin D supplementation is necessary.

Formatted
Numbering

| 149) Sufficient skin exposure to solar UV radiation of the appropriate wavelength is essential for maintaining adequate vitamin D status in the UK. A clear recommendation is required on the length and intensity of exposure necessary to balance maintenance of vitamin D status with the risk of developing skin cancer.

Formatted
Numbering

| 150) There is a need to understand better the effects of adiposity on 25-hydroxyvitamin D concentration and its implications for vitamin D requirements.

Formatted
Numbering

| 151) Vulnerable people include infants and young children (up to the age of 3 years), older adults, pregnant and breastfeeding women, particularly but not exclusively those from - ethnic minority backgrounds. Dietary supplements of vitamin D remain necessary to safeguard these groups against deficiency and its adverse effects on health.

Formatted
Numbering

References

- Agarwal KS, Mughal MZ, Upadhyay P, Berry JL, Mawer EB & Puliyeel JM (2002) The impact of atmospheric pollution on vitamin D status of infants and toddlers in Delhi, India. *Arch Dis Child* **87**, 111-113.
- Ahmed I, Atiq M, Iqbal J, Khurshid M & Whittaker P (1995) Vitamin D deficiency rickets in breast-fed infants presenting with hypocalcaemic seizures. *Acta Paediatr* **84**, 941-942.
- Ahonen MH, Tenkanen L, Teppo L, Hakama M & Tuohimaa P (2000) Prostate cancer risk and prediagnostic serum 25-hydroxyvitamin D levels (Finland). *Cancer Causes Control* **11**, 847-852.
- Ala-Houhala M (1985) 25-Hydroxyvitamin D levels during breast-feeding with or without maternal or infantile supplementation of vitamin D. *J Pediatr Gastroenterol Nutr* **4**, 220-226.
- Ala-Houhala M, Koskinen T, Terho A, Koivula T & Visakorpi J (1986) Maternal compared with infant vitamin D supplementation. *Arch Dis Child* **61**, 1159-1163.
- Allgrove J (2004) Is nutritional rickets returning? *Arch Dis Child* **89**, 699-701.
- Armas LA, Hollis BW & Heaney RP (2004) Vitamin D₂ is much less effective than vitamin D₃ in humans. *J Clin Endocrinol Metab* **89**, 5387-5391.
- Arunabh S, Pollack S, Yeh J & Aloia JF (2003) Body Fat Content and 25-Hydroxyvitamin D Levels in Healthy Women. *J Clin Endocrinol Metab* **88**, 157-161.
- Ashraf S & Mughal MZ (2002) The prevalence of rickets among non-Caucasian children. *Arch Dis Child* **87**, 263-26a.
- Avenell, A., Gillespie, W.J., Gillespie, L.D. & O'Connell, D.L. (2006) Vitamin D and vitamin D analogues for preventing fractures associated with involutional and post-menopausal osteoporosis (Cochrane Review). *The Cochrane Database of Systematic Reviews*, Issue 3.
- Awumey EM, Mitra DA, Hollis BW, Kumar R & Bell NH (1998) Vitamin D metabolism is altered in Asian Indians in the southern United States: a clinical research center study. *J Clin Endocrinol Metab* **83**, 169-173.
- Barnes PF, Modlin RL, Bikle DD & Adams JS (1989) Transpleural gradient of 1,25-dihydroxyvitamin D in tuberculous pleuritis. *J Clin Invest* **83**, 1527-1532.
- Bates CJ, Carter GD, Mishra GD, O'Shea D, Jones J & Prentice A (2003) In a population study, can parathyroid hormone aid the definition of adequate vitamin D status? A study of people aged 65 years and over from the British National Diet and Nutrition Survey. *Osteoporos Int* **14**, 152-159.
- Bell NH, Epstein S, Greene A, Shary J, Oexmann MJ & Shaw S (1985) Evidence for alteration of the vitamin D-endocrine system in obese subjects. *J Clin Invest* **76**, 370-373.

- Bertone-Johnson ER, Chen WY, Holick MF, Hollis BW, Colditz GA, Willett WC & Hankinson SE (2005) Plasma 25-Hydroxyvitamin D and 1,25-Dihydroxyvitamin D and Risk of Breast Cancer. *Cancer Epidemiol Biomarkers Prev* **14**, 1991-1997.
- Bikle DD, Gee E, Halloran B, Kowalski MA, Ryzen E & Haddad JG (1986) Assessment of the free fraction of 25-hydroxyvitamin D in serum and its regulation by albumin and the vitamin D-binding protein. *J Clin Endocrinol Metab* **63**, 954-959.
- Bikle DD, Siiteri PK, Ryzen E & Haddad JG (1985) Serum protein binding of 1,25-dihydroxyvitamin D: a re-evaluation by direct measurement of free metabolite levels. *J Clin Endocrinol Metab* **61**, 969-975.
- Binkley N, Krueger D, Cowgill CS, Plum L, Lake E, Hansen KE, DeLuca HF & Drezner MK (2004) Assay variation confounds the diagnosis of hypovitaminosis D: a call for standardization. *J Clin Endocrinol Metab* **89**, 3152-3157.
- Bischoff-Ferrari HA, Dawson-Hughes B, Willett WC, Staehelin HB, Bazemore MG, Zee RY & Wong JB (2004) Effect of Vitamin D on falls: a meta-analysis. *JAMA* **291**, 1999-2006.
- Bischoff-Ferrari HA, Willett WC, Wong JB, Giovannucci E, Dietrich T & Dawson-Hughes B (2005) Fracture prevention with vitamin D supplementation: a meta-analysis of randomized controlled trials. *JAMA* **293**, 2257-2264.
- Bischoff-Ferrari HA, Giovannucci E, Willett WC, Dietrich T & Dawson-Hughes B (2006) Estimation of optimal serum concentrations of 25-hydroxyvitamin D for multiple health outcomes. *Am J Clin Nutr* **84**, 18-28.
- Blank S, Scanlon KS, Sinks TH, Lett S & Falk H (1995) An outbreak of hypervitaminosis D associated with the overfortification of milk from a home-delivery dairy. *Am J Public Health* **85**, 656-659.
- Borissova AM, Tankova T, Kirilov G, Dakovska L & Kovacheva R (2003) The effect of vitamin D₃ on insulin secretion and peripheral insulin sensitivity in type 2 diabetic patients. *Int J Clin Pract* **57**, 258-261.
- Bostick RM, Potter JD, Sellers TA, McKenzie DR, Kushi LH & Folsom AR (1993) Relation of calcium, vitamin D, and dairy food intake to incidence of colon cancer among older women. The Iowa Women's Health Study. *Am J Epidemiol* **137**, 1302-1317.
- Boucher BJ (2006) Hypovitaminosis D and risk of Type 2 diabetes in British South Asians. *Diabet Med* **23**, 336.
- Braun MM, Helzlsouer KJ, Hollis BW & Comstock GW (1995) Prostate cancer and prediagnostic levels of serum vitamin D metabolites (Maryland, United States). *Cancer Causes Control* **6**, 235-239.
- British Pediatric Association (1956) Hypercalcemia in infants and vitamin D. *BMJ* **2**, 149-155.
- Buffington C, Walker B, Cowan GS, Jr. & Scruggs D (1993) Vitamin D Deficiency in the

- Morbidly Obese. *Obes Surg* **3**, 421-424.
- Callaghan AL, Moy RJ, Booth IW, DeBelle GD & Shaw NJ (2006) Incidence of symptomatic vitamin D deficiency. *Arch Dis Child* **91**, 606-607.
- Carter GD, Carter R, Jones J & Berry J (2004) How Accurate Are Assays for 25-Hydroxyvitamin D? Data from the International Vitamin D External Quality Assessment Scheme. *Clin Chem* **50**, 2195-2197.
- Chief Medical Officer. (2005) Meeting the need for vitamin D. *CMO Update* **42**, 6.
- Chiu KC, Chu A, Go VL & Saad MF (2004) Hypovitaminosis D is associated with insulin resistance and β cell dysfunction. *Am J Clin Nutr* **79**, 820-825.
- Compston JE, Vedi S, Ledger JE, Webb A, Gazet JC & Pilkington TR (1981) Vitamin D status and bone histomorphometry in gross obesity. *Am J Clin Nutr* **34**, 2359-2363.
- Corder EH, Guess HA, Hulka BS, Friedman GD, Sadler M, Vollmer RT, Lobaugh B, Drezner MK, Vogelmann JH & Orentreich N (1993) Vitamin D and prostate cancer: a prediagnostic study with stored sera. *Cancer Epidemiol Biomarkers Prev* **2**, 467-472.
- Crocombe S, Mughal MZ & Berry JL (2004) Symptomatic vitamin D deficiency among non-Caucasian adolescents living in the United Kingdom. *Arch Dis Child* **89**, 197-19a.
- CRUK (2006) Our position on vitamin D. <http://cancerresearchuk.org/sunsmart/forprofessionals/vitaminD/>.
- Cui Y & Rohan TE (2006) Vitamin D, Calcium, and Breast Cancer Risk: A Review. *Cancer Epidemiol Biomarkers Prev* **15**, 1427-1437.
- d'Alesio A, Garabedian M, Sabatier JP, Guaydier-Souquieres G, Marcelli C, Lemacon A, Walrant-Debray O & Jehan F (2005) Two single-nucleotide polymorphisms in the human vitamin D receptor promoter change protein-DNA complex formation and are associated with height and vitamin D status in adolescent girls. *Hum Mol Genet* **14**, 3539-3548.
- Das G, Crocombe S, McGrath M, Berry J & Mughal Z (2006) Hypovitaminosis D among healthy adolescent girls attending an inner city school. *Arch Dis Child* **91**, 569-572.
- Datta S, Alfaham M, Davies DP, Dunstan F, Woodhead S, Evans J & Richards B (2002) Vitamin D deficiency in pregnant women from a non-European ethnic minority population--an interventional study. *BJOG* **109**, 905-908.
- Davies PS, Bates CJ, Cole TJ, Prentice A & Clarke PC (1999) Vitamin D: seasonal and regional differences in preschool children in Great Britain. *Eur J Clin Nutr* **53**, 195-198.
- Dawson-Hughes B, Heaney RP, Holick MF, Lips P, Meunier PJ & Vieth R (2005) Estimates of optimal vitamin D status. *Osteoporos Int* **16**, 713-716.
- DeLuca HF (2004) Overview of general physiologic features and functions of vitamin D. *Am*

J Clin Nutr **80**, 1689S-1696.

Department of Health (1991a) *Dietary Reference Values for Food Energy and Nutrients for the United Kingdom* no. 41. London: HMSO.

Department of Health (1991b) *The fortification of yellow fats with vitamins A and D* no. 40. London: HMSO.

Department of Health (1998) *Nutrition and Bone Health: with particular reference to calcium and vitamin D* no. 49. London: The Stationary Office.

Department of Health (2002) *Scientific Review of the Welfare Food Scheme* no. 51. London: TSO.

Department of Health and Social Security (1980) *Rickets and Osteomalacia* no. 19. London: HMSO.

Dunnigan MG, Glekin BM, Henderson JB, McIntosh WB, Sumner D & Sutherland GR (1985) Prevention of rickets in Asian children: assessment of the Glasgow campaign. *Br Med J (Clin Res Ed)* **291**, 239-242.

Dunnigan MG, Henderson JB, Hole DJ, Barbara ME & Berry JL (2005) Meat consumption reduces the risk of nutritional rickets and osteomalacia. *Br J Nutr* **94**, 983-991.

el Sonbaty MR & Abdul-Ghaffar NU (1996) Vitamin D deficiency in veiled Kuwaiti women. *Eur J Clin Nutr* **50**, 315-318.

EURODIAB Substudy 2 Study Group (1999) Vitamin D supplement in early childhood and risk for Type I (insulin-dependent) diabetes mellitus. *Diabetologia* **42**, 51-54.

Expert Group on Vitamins and Minerals (2003) *Safe Upper Levels for Vitamins and Minerals*. London: FSA.

Farrerons J, Barnadas M, Rodriguez J, Renau A, Yoldi B, Lopez-Navidad A & Moragas J (1998) Clinically prescribed sunscreen (sun protection factor 15) does not decrease serum vitamin D concentration sufficiently either to induce changes in parathyroid function or in metabolic markers. *Br J Dermatol* **139**, 422-427.

Feskanich D, Ma J, Fuchs CS, Kirkner GJ, Hankinson SE, Hollis BW & Giovannucci EL (2004) Plasma vitamin D metabolites and risk of colorectal cancer in women. *Cancer Epidemiol Biomarkers Prev* **13**, 1502-1508.

Fleet JC (2004) Genomic and proteomic approaches for probing the role of vitamin D in health. *Am J Clin Nutr* **80**, 1730S-1734.

Finch S, Doyle W, Lowe C, Bates CJ, Prentice A, Smithers G & Clarke CC (1998) *National Diet and Nutrition Survey: people aged 65 years and over*. London, The Stationary Office

Forman JP, Bischoff-Ferrari HA, Willett WC, Stampfer MJ & Curhan GC (2005) Vitamin D Intake and Risk of Incident Hypertension: Results From Three Large Prospective

- Cohort Studies. *Hypertension* **46**, 676-682.
- Fraser, DR (1967) The relation between infantile hypercalcemia and vitamin D--public health implications in North America. *Pediatrics* **40**(6), 1050-1061
- Gann PH, Ma J, Hennekens CH, Hollis BW, Haddad JG & Stampfer MJ (1996) Circulating vitamin D metabolites in relation to subsequent development of prostate cancer. *Cancer Epidemiol Biomarkers Prev* **5**, 121-126.
- Gannage-Yared MH, CHEMALI RANA, YAACOUB NAJW & HALABY GEOR (2000) Hypovitaminosis D in a Sunny Country: Relation to Lifestyle and Bone Markers. *Journal of Bone and Mineral Research* **15**, 1856-1862.
- Garland C, Shekelle RB, Barrett-Connor E, Criqui MH, Rossof AH & Paul O (1985) Dietary vitamin D and calcium and risk of colorectal cancer: a 19-year prospective study in men. *Lancet* **1**, 307-309.
- Garland CF, Comstock GW, Garland FC, Helsing KJ, Shaw EK & Gorham ED (1989) Serum 25-hydroxyvitamin D and colon cancer: eight-year prospective study. *Lancet* **2**, 1176-1178.
- Garland CF, Garland FC, Gorham ED, Lipkin M, Newmark H, Mohr SB & Holick MF (2006) The role of vitamin D in cancer prevention. *Am J Public Health* **96**, 252-261.
- Ghannam NN, Hammami MM, Bakheet SM & Khan BA (1999) Bone mineral density of the spine and femur in healthy Saudi females: relation to vitamin D status, pregnancy, and lactation. *Calcif Tissue Int* **65**, 23-28.
- Giovannucci E (2006) The epidemiology of vitamin D and colorectal cancer: recent findings. *Curr Opin Gastroenterol* **22**, 24-29.
- Giovannucci E (2005) The epidemiology of vitamin D and cancer incidence and mortality: A review (United States). *Cancer Causes and Control* **16**, 83-95.
- Giovannucci E, Liu Y, Rimm EB, Hollis BW, Fuchs CS, Stampfer MJ & Willett WC (2006) Prospective Study of Predictors of Vitamin D Status and Cancer Incidence and Mortality in Men. *J Natl Cancer Inst* **98**, 451-459.
- Gorham ED, Garland CF, Garland FC, Grant WB, Mohr SB, Lipkin M, Newmark HL, Giovannucci E, Wei M & Holick MF (2005) Vitamin D and prevention of colorectal cancer. *J Steroid Biochem Mol Biol* **97**, 179-194.
- Grant AM, Avenell A, Campbell MK, *et al.* (2005) Oral vitamin D3 and calcium for secondary prevention of low-trauma fractures in elderly people (Randomized Evaluation of Calcium Or vitamin D, RECORD): a randomized placebo-controlled trial. *Lancet* **365**, 1621-1628.
- Greer FR, Hollis BW, Cripps DJ & Tsang RC (1984) Effects of maternal ultraviolet B irradiation on vitamin D content of human milk. *J Pediatr* **105**, 431-433.

- Gregory J, Lowe S, Bates CJ, Prentice A, Jackson LV, Smithers G & Wenlock R (2000) *National Diet and Nutrition Survey: young people aged 4 to 18 years*. London: The Stationary Office.
- Gregory JR, Collins DL, Davies PSW, Hughes JM & Clarke PC (1995) *National Diet and Nutrition Survey: children aged 1½-4½ years. Volume 1: report of the diet and nutrition survey*. London: HMSO.
- Haddad JG (1995) Plasma vitamin D-binding protein (Gc-globulin): multiple tasks. *J Steroid Biochem Mol Biol* **53**, 579-582.
- Hanley DA & Davison KS (2005) Vitamin D Insufficiency in North America. *J Nutr* **135**, 332-337.
- Harris SS & Dawson-Hughes B (1998) Seasonal changes in plasma 25-hydroxyvitamin D concentrations of young American black and white women. *Am J Clin Nutr* **67**, 1232-1236.
- Hathcock JN, Shao A, Vieth R, Heaney R (2007) Risk assessment for vitamin D. *Am J Clin Nutr* **85**, 6-18
- Heaney RP (2004) Functional indices of vitamin D status and ramifications of vitamin D deficiency. *Am J Clin Nutr* **80**, 1706S-1709.
- Heaney RP, Davies KM, Chen TC, Holick MF & Barger-Lux MJ (2003) Human serum 25-hydroxycholecalciferol response to extended oral dosing with cholecalciferol. *Am J Clin Nutr* **77**, 204-210.
- Henderson L, Irving K & Gregory J (2003) *The National Diet and Nutrition Survey: adults aged 19 to 64 years. Volume 3: Vitamin and mineral intake and urinary analytes*. London, TSO.
- Hey H, Stokholm KH, Lund B, Lund B & Sorensen OH (1982) Vitamin D deficiency in obese patients and changes in circulating vitamin D metabolites following jejunoileal bypass. *Int J Obes* **6**, 473-479.
- Hirani V & Primatesta P (2005) Vitamin D concentrations among people aged 65 years and over living in private households and institutions in England: population survey. *Age Ageing* **34**, 485-491.
- Holick MF, Matsuoka LY & Wortsman J (1995) Regular use of sunscreen on vitamin D levels. *Arch Dermatol* **131**, 1337-1339.
- Holick MF (2004) Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *Am J Clin Nutr* **80**, 1678S-1688.
- Holick MF (2006) Resurrection of vitamin D deficiency and rickets. *J Clin Invest* **116**, 2062-2072.
- Hollis BW (2005) Circulating 25-hydroxyvitamin D levels indicative of vitamin D sufficiency: implications for establishing a new effective dietary intake

- recommendation for vitamin D. *J Nutr* **135**, 317-322.
- Hollis BW & Wagner CL (2004) Vitamin D requirements during lactation: high-dose maternal supplementation as therapy to prevent hypovitaminosis D for both the mother and the nursing infant. *Am J Clin Nutr* **80**, 1752S-1758.
- Houghton LA & Vieth R (2006) The case against ergocalciferol (vitamin D₂) as a vitamin supplement. *Am J Clin Nutr* **84**, 694-697.
- Hyldstrup L, Andersen T, McNair P, Breum L & Transbol I (1993) Bone metabolism in obesity: changes related to severe overweight and dietary weight reduction. *Acta Endocrinol (Copenh)* **129**, 393-398.
- Hyponen E, Laara E, Reunanen A, Jarvelin MR & Virtanen SM (2001) Intake of vitamin D and risk of type 1 diabetes: a birth-cohort study. *Lancet* **358**, 1500-1503.
- Hyponen E, Sovio U, Wjst M, Patel S, Pekkanen J, Hartikainen AL & Jarvelinb MR (2004) Infant vitamin d supplementation and allergic conditions in adulthood: northern Finland birth cohort 1966. *Ann N Y Acad Sci* **1037**, 84-95.
- Hyponen E & Power C (2006) Vitamin D Status and Glucose Homeostasis in the 1958 British Birth Cohort: The role of obesity. *Diabetes Care* **29**, 2244-2246.
- Institute of Medicine (1997) *Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride*. Washington, D.C.: National Academy Press.
- Iqbal J, Walker C & Swift P (2001) The continuing problem of vitamin D deficiency in pregnant Asian women and their offspring; an interface audit as a prelude to action. *Arch Dis Child* **84**, A10-A68.
- Jackson RD, LaCroix AZ, Gass M, *et al.* (2006) Calcium plus vitamin D supplementation and the risk of fractures. *N Engl J Med* **354**, 669-683.
- Jacobus CH, Holick MF, Shao Q, Chen TC, Holm IA, Kolodny JM, Fuleihan GE & Seely EW (1992) Hypervitaminosis D associated with drinking milk. *N Engl J Med* **326**, 1173-1177.
- Jarvinen R, Knekt P, Hakulinen T & Aromaa A (2001) Prospective study on milk products, calcium and cancers of the colon and rectum. *Eur J Clin Nutr* **55**, 1000-1007.
- Javaid MK, Crozier SR, Harvey NC, Gale CR, Dennison EM, Boucher BJ, Arden NK, Godfrey KM & Cooper C (2006) Maternal vitamin D status during pregnancy and childhood bone mass at age 9 years: a longitudinal study. *Lancet* **367**, 36-43.
- John EM, Schwartz GG, Koo J, Van Den Berg D & Ingles SA (2005) Sun Exposure, Vitamin D Receptor Gene Polymorphisms, and Risk of Advanced Prostate Cancer. *Cancer Res* **65**, 5470-5479.
- Kearney J, Giovannucci E, Rimm EB, Ascherio A, Stampfer MJ, Colditz GA, Wing A, Kampman E & Willett WC (1996) Calcium, vitamin D, and dairy foods and the occurrence of colon cancer in men. *Am J Epidemiol* **143**, 907-917.

- Kreiter SR, Schwartz RP, Kirkman HN, Jr., Charlton PA, Calikoglu AS & Davenport ML (2000) Nutritional rickets in African American breast-fed infants. *J Pediatr* **137**, 153-157.
- Laaksi IT, Ruohola J-PS, Ylikomi TJ, Auvinen A, Haataja RI, Pihlajamaki HK & Tuohimaa PJ (2006) Vitamin D fortification as public health policy: significant improvement in vitamin D status in young Finnish men. *Eur J Clin Nutr* **60**, 1035-1038.
- Laaksonen M, Karkkainen M, Outila T, Vanninen T, Ray C & Lamberg-Allardt C (2002) Vitamin D receptor gene BsmI-polymorphism in Finnish premenopausal and postmenopausal women: its association with bone mineral density, markers of bone turnover, and intestinal calcium absorption, with adjustment for lifestyle factors. *J Bone Miner Metab* **20**, 383-390.
- Ladhani S, Srinivasan L, Buchanan C & Allgrove J (2004) Presentation of vitamin D deficiency. *Arch Dis Child* **89**, 781-784.
- Lane NE, Gore LR, Cummings SR, Hochberg MC, Scott JC, Williams EN & Nevitt MC (1999) Serum vitamin D levels and incident changes of radiographic hip osteoarthritis: a longitudinal study. Study of Osteoporotic Fractures Research Group. *Arthritis Rheum* **42**, 854-860.
- Lauridsen AL, Vestergaard P, Hermann AP, Brot C, Heickendorff L, Mosekilde L & Nexø E (2005) Plasma concentrations of 25-hydroxy-vitamin D and 1,25-dihydroxy-vitamin D are related to the phenotype of Gc (vitamin D-binding protein): a cross-sectional study on 595 early postmenopausal women. *Calcif Tissue Int* **77**, 15-22.
- Lawson DE, Paul AA, Black AE, Cole TJ, Mandal AR & Davie M (1979) Relative contributions of diet and sunlight to vitamin D state in the elderly. *Br Med J* **2**, 303-305.
- Lawson M & Thomas M (1999) Vitamin D concentrations in Asian children aged 2 years living in England: population survey. *BMJ* **318**, 28.
- Lawson M, Thomas M & Hardiman A (1999) Dietary and lifestyle factors affecting plasma vitamin D levels in Asian children living in England. *Eur J Clin Nutr* **53**, 268-272.
- Lensmeyer GL, Wiebe DA, Binkley N & Drezner MK (2006) HPLC method for 25-hydroxyvitamin D measurement: comparison with contemporary assays. *Clin Chem* **52**, 1120-1126.
- Leventis P, Garrison L, Sibley M, Peterson P, Egerton M, Levin G & Kiely P (2005) Underestimation of Serum 25-Hydroxyvitamin D by the Nichols Advantage Assay in Patients Receiving Vitamin D Replacement Therapy. *Clin Chem* **51**, 1072-1074.
- Liel Y, Ulmer E, Shary J, Hollis BW & Bell NH (1988) Low circulating vitamin D in obesity. *Calcif Tissue Int* **43**, 199-201.
- Lin J, Zhang SM, Cook NR, Manson JE, Lee IM & Buring JE (2005) Intakes of Calcium and Vitamin D and Risk of Colorectal Cancer in Women. *Am J Epidemiol* **161**, 755-764.

- Lind L, Lithell H, Skarfors E, Wide L & Ljunghall S (1988a) Reduction of blood pressure by treatment with alphacalcidol. A double-blind, placebo-controlled study in subjects with impaired glucose tolerance. *Acta Med Scand* **223**, 211-217.
- Lind L, Wengle B, Wide L, Sorensen OH & Ljunghall S (1988b) Hypertension in primary hyperparathyroidism--reduction of blood pressure by long-term treatment with vitamin D (alphacalcidol). A double-blind, placebo-controlled study. *Am J Hypertens* **1**, 397-402.
- Lips P (2004) Which circulating level of 25-hydroxyvitamin D is appropriate? *J Steroid Biochem Mol Biol* **89-90**, 611-614.
- Liu PT, Stenger S, Li H, *et al.* (2006) Toll-Like Receptor Triggering of a Vitamin D-Mediated Human Antimicrobial Response. *Science* **311**, 1770-1773.
- Lo CW, Paris PW & Holick MF (1986) Indian and Pakistani immigrants have the same capacity as Caucasians to produce vitamin D in response to ultraviolet irradiation. *Am J Clin Nutr* **44**, 683-685.
- Looker AC (2005) Body Fat and Vitamin D Status in Black Versus White Women. *J Clin Endocrinol Metab* **90**, 635-640.
- Lowe LC, Guy M, Mansi JL, Peckitt C, Bliss J, Wilson RG & Colston KW (2005) Plasma 25-hydroxy vitamin D concentrations, vitamin D receptor genotype and breast cancer risk in a UK Caucasian population. *Eur J Cancer* **41**, 1164-1169.
- Lund B, Sorensen OH, Lund B, Bishop JE & Norman AW (1980) Vitamin D metabolism in hypoparathyroidism. *J Clin Endocrinol Metab* **51**, 606-610.
- Marks R, Foley PA, Jolley D, Knight KR, Harrison J & Thompson SC (1995) The effect of regular sunscreen use on vitamin D levels in an Australian population. Results of a randomized controlled trial. *Arch Dermatol* **131**, 415-421.
- Martinez ME, Giovannucci EL, Colditz GA, Stampfer MJ, Hunter DJ, Speizer FE, Wing A & Willett WC (1996) Calcium, vitamin D, and the occurrence of colorectal cancer among women. *J Natl Cancer Inst* **88**, 1375-1382.
- Matsuoka LY, Ide L, Wortsman J, MacLaughlin JA & Holick MF (1987) Sunscreens suppress cutaneous vitamin D₃ synthesis. *J Clin Endocrinol Metab* **64**, 1165-1168.
- Matsuoka LY, Wortsman J, Dannenberg MJ, Hollis BW, Lu Z & Holick MF (1992) Clothing prevents ultraviolet-B radiation-dependent photosynthesis of vitamin D₃. *J Clin Endocrinol Metab* **75**, 1099-1103.
- Matsuoka LY, Wortsman J, Haddad JG, Kolm P & Hollis BW (1991) Racial pigmentation and the cutaneous synthesis of vitamin D. *Arch Dermatol* **127**, 536-538.
- Matsuoka LY, Wortsman J, Hanifan N & Holick MF (1988) Chronic sunscreen use decreases circulating concentrations of 25-hydroxyvitamin D. A preliminary study. *Arch Dermatol* **124**, 1802-1804.

- Mawer EB, Backhouse J, Holman CA, Lumb GA & Stanbury SW (1972) The distribution and storage of vitamin D and its metabolites in human tissues. *Clin Sci* **43**, 413-431.
- McAlindon TE, Felson DT, Zhang Y, Hannan MT, Aliabadi P, Weissman B, Rush D, Wilson PWF & Jacques P (1996) Relation of Dietary Intake and Serum Levels of Vitamin D to Progression of Osteoarthritis of the Knee among Participants in the Framingham Study. *Ann Intern Med* **125**, 353-359.
- McCullough ML, Robertson AS, Rodriguez C, Jacobs EJ, Chao A, Carolyn J, Calle EE, Willett WC & Thun MJ (2003) Calcium, vitamin D, dairy products, and risk of colorectal cancer in the Cancer Prevention Study II Nutrition Cohort (United States). *Cancer Causes Control* **14**, 1-12.
- McKinlay, A. (2006) Workshop Round-up Session Rapporteur's Report. *Progress in Biophysics and Molecular Biology* **92**, 179-184
- Mehta, DK (2006) *British National Formulary, 52nd Edition*. Pharmaceutical Press, UK.
- Ministry of Health and Department of Health for Scotland (1957) *Report of the joint sub-committee on welfare food*. London: HMSO.
- Mughal MZ, Salama H, Greenaway T, Laing I & Mawer EB (1999) Lesson of the week: Florid rickets associated with prolonged breast feeding without vitamin D supplementation. *BMJ* **318**, 39-40.
- Munger KL, Levin LI, Hollis BW, Howard NS & Ascherio A (2006) Serum 25-Hydroxyvitamin D Levels and Risk of Multiple Sclerosis. *JAMA* **296**, 2832-2838.
- Munger KL, Zhang SM, O'Reilly E, Hernan MA, Olek MJ, Willett WC & Ascherio A (2004) Vitamin D intake and incidence of multiple sclerosis. *Neurology* **62**, 60-65.
- National Institute for Health and Clinical Excellence (2003) Antenatal care. Routine care for the healthy pregnant woman.
- Need AG, Morris HA, Horowitz M & Nordin C (1993) Effects of skin thickness, age, body fat, and sunlight on serum 25-hydroxyvitamin D. *Am J Clin Nutr* **58**, 882-885.
- Ness AR, Frankel SJ, Gunnell DJ & Smith GD (1999) Are we really dying for a tan? *BMJ* **319**, 114-116.
- Nomura AM, Stemmermann GN, Lee J, Kolonel LN, Chen TC, Turner A & Holick MF (1998) Serum vitamin D metabolite levels and the subsequent development of prostate cancer (Hawaii, United States). *Cancer Causes Control* **9**, 425-432.
- Norman AW (1998) Sunlight, season, skin pigmentation, vitamin D, and 25-hydroxyvitamin D: integral components of the vitamin D endocrine system. *Am J Clin Nutr* **67**, 1108-1110.
- Odeka E & Tan J (2005) Nutritional rickets is increasingly diagnosed in children of ethnic origin. *Arch Dis Child* **90**, 1203-1204.

- Ogunkolade BW, Boucher BJ, Prah JM, *et al.* (2002) Vitamin D receptor (VDR) mRNA and VDR protein levels in relation to vitamin D status, insulin secretory capacity, and VDR genotype in Bangladeshi Asians. *Diabetes* **51**, 2294-2300.
- Pal BR, Marshall T, James C & Shaw NJ (2003) Distribution analysis of vitamin D highlights differences in population subgroups: preliminary observations from a pilot study in UK adults. *J Endocrinol* **179**, 119-129.
- Pal BR & Shaw NJ (2001) Rickets resurgence in the United Kingdom: improving antenatal management in Asians. *J Pediatr* **139**, 337-338.
- Parikh SJ, Edelman M, Uwaifo GI, Freedman RJ, Semega-Janneh M, Reynolds J & Yanovski JA (2004) The Relationship between Obesity and Serum 1,25-Dihydroxy Vitamin D Concentrations in Healthy Adults. *J Clin Endocrinol Metab* **89**, 1196-1199.
- Pettifor JM, Bikle DD, Cavaleros M, Zachen D, Kamdar MC & Ross FP (1995) Serum Levels of Free 1,25-Dihydroxyvitamin D in Vitamin D Toxicity. *Ann Intern Med* **122**, 511-513.
- Pfeifer M, Begerow B, Minne HW, Nachtigall D & Hansen C (2001) Effects of a Short-Term Vitamin D3 and Calcium Supplementation on Blood Pressure and Parathyroid Hormone Levels in Elderly Women. *J Clin Endocrinol Metab* **86**, 1633-1637.
- Platz EA, Leitzmann MF, Hollis BW, Willett WC & Giovannucci E (2004) Plasma 1,25-dihydroxy- and 25-hydroxyvitamin D and subsequent risk of prostate cancer. *Cancer Causes Control* **15**, 255-265.
- Porthouse J, Cockayne S, King C, *et al.* (2005) Randomized controlled trial of calcium and supplementation with cholecalciferol (vitamin D3) for prevention of fractures in primary care. *BMJ* **330**, 1003.
- Ren S, Nguyen L, Wu S, Encinas C, Adams JS & Hewison M (2005) Alternative Splicing of Vitamin D-24-Hydroxylase: A NOVEL MECHANISM FOR THE REGULATION OF EXTRARENAL 1,25-DIHYDROXYVITAMIN D SYNTHESIS. *J Biol Chem* **280**, 20604-20611.
- Robinson PD, Hogler W, Craig ME, Verge CF, Walker JL, Piper AC, Woodhead HJ, Cowell CT & Ambler GR (2006) The re-emerging burden of rickets: A decade of experience from Sydney. *Arch Dis Child* **91**, 564-568.
- Ruston D, Henderson L, Gregory J, Bates CJ, Prentice A, Birch M, Swan G & Farron M (2004) *The National Diet and Nutrition Survey: adults aged 19 to 64 years. Volume 4: Nutritional Status (anthropometry and blood analytes), blood pressure and physical activity*. London: TSO.
- Schleithoff SS, Zittermann A, Tenderich G, Berthold HK, Stehle P & Koerfer R (2006) Vitamin D supplementation improves cytokine profiles in patients with congestive heart failure: a double-blind, randomized, placebo-controlled trial. *Am J Clin Nutr* **83**, 754-759.
- Schwartz GG, Eads D, Rao A, *et al.* (2004) Pancreatic cancer cells express 25-

hydroxyvitamin D-1{alpha}-hydroxylase and their proliferation is inhibited by the prohormone 25-hydroxyvitamin D3. *Carcinogenesis* **25**, 1015-1026.

Scientific Advisory Committee on Nutrition (2005) *Review of Dietary Advice on Vitamin A*. London, TSO.

Scientific Committee on Food & Scientific Panel on Dietetic Products NaA (2006) Tolerable upper intake levels for vitamins and minerals. <http://www.efsa.europa.eu/en/science/nda.html>.

Scragg R, Khaw KT & Murphy S (1995) Effect of winter oral vitamin D3 supplementation on cardiovascular risk factors in elderly adults. *Eur J Clin Nutr* **49**, 640-646.

Segersten U, Correa P, Hewison M, Hellman P, Dralle H, Carling T, Akerstrom G & Westin G (2002) 25-Hydroxyvitamin D3-1{alpha}-Hydroxylase Expression in Normal and Pathological Parathyroid Glands. *J Clin Endocrinol Metab* **87**, 2967-2972.

Shaw NJ & Pal BR (2002) Vitamin D deficiency in UK Asian families: activating a new concern. *Arch Dis Child* **86**, 147-149.

Shenoy SD, Swift P, Cody D & Iqbal J (2005) Maternal vitamin D deficiency, refractory neonatal hypocalcaemia, and nutritional rickets. *Arch Dis Child* **90**, 437-438.

Skinner HG, Michaud DS, Giovannucci E, Willett WC, Colditz GA & Fuchs CS (2006) Vitamin D Intake and the Risk for Pancreatic Cancer in Two Cohort Studies. *Cancer Epidemiol Biomarkers Prev* **15**, 1688-1695.

Snijder MB, van Dam RM, Visser M, Deeg DJ, Dekker JM, Bouter LM, Seidell JC & Lips P (2005) Adiposity in relation to vitamin D status and parathyroid hormone levels: a population-based study in older men and women. *J Clin Endocrinol Metab* **90**, 4119-4123.

Specker BL (1994) Do North American women need supplemental vitamin D during pregnancy or lactation? *Am J Clin Nutr* **59**, 484S-490.

Specker BL, Valanis B, Hertzberg V, Edwards N & Tsang RC (1985) Sunshine exposure and serum 25-hydroxyvitamin D concentrations in exclusively breast-fed infants. *J Pediatr* **107**, 372-376.

Stolzenberg-Solomon RZ, Vieth R, Azad A, Pietinen P, Taylor PR, Virtamo J & Albanes D (2006) A Prospective Nested Case-Control Study of Vitamin D Status and Pancreatic Cancer Risk in Male Smokers. *Cancer Res* **66**, 10213-10219.

Teegarden D, Meredith SC & Sitrin MD (1991) Determination of the affinity of vitamin D metabolites to serum vitamin D binding protein using assay employing lipid-coated polystyrene beads. *Anal Biochem* **199**, 293-299.

Thacher TD, Fischer PR, Pettifor JM, Lawson JO, Isichei CO, Reading JC & Chan GM (1999) A Comparison of Calcium, Vitamin D, or Both for Nutritional Rickets in Nigerian Children. *N Engl J Med* **341**, 563-568.

- Timms PM, Mannan N, Hitman GA, Noonan K, Mills PG, Syndercombe-Court, Aganna E, Price CP & Boucher BJ (2002) Circulating MMP9, vitamin D and variation in the TIMP-1 response with VDR genotype: mechanisms for inflammatory damage in chronic disorders? *QJM* **95**, 787-796.
- Tjellesen L, Hummer L, Christiansen C & Rodbro P (1986) Serum concentration of vitamin D metabolites during treatment with vitamin D2 and D3 in normal premenopausal women. *Bone Miner* **1**, 407-413.
- Trang HM, Cole DE, Rubin LA, Pierratos A, Siu S & Vieth R (1998) Evidence that vitamin D3 increases serum 25-hydroxyvitamin D more efficiently than does vitamin D2. *Am J Clin Nutr* **68**, 854-858.
- Tuohimaa P, Tenkanen L, Ahonen M, *et al.* (2004) Both high and low levels of blood vitamin D are associated with a higher prostate cancer risk: a longitudinal, nested case-control study in the Nordic countries. *Int J Cancer* **108**, 104-108.
- Ustianowski A, Shaffer R, Collin S, Wilkinson RJ & Davidson RN (2005) Prevalence and associations of vitamin D deficiency in foreign-born persons with tuberculosis in London. *J Infect* **50**, 432-437.
- Valimaki VV, Loyttyneimi E & Valimaki MJ (2006) Vitamin D fortification of milk products does not resolve hypovitaminosis D in young Finnish men. *Eur J Clin Nutr*. Epub ahead of print
- van Driel M, Koedam M, Buurman CJ, Hewison M, Chiba H, Uitterlinden AG, Pols HAP & van Leeuwen JPTM (2006) Evidence for auto/paracrine actions of vitamin D in bone: 1{alpha}-hydroxylase expression and activity in human bone cells. *FASEB J* **20**, 2417-2419.
- Vieth R (1990) The mechanisms of vitamin D toxicity. *Bone Miner* **11**, 267-272.
- Vieth R (1999) Vitamin D supplementation, 25-hydroxyvitamin D concentrations, and safety. *Am J Clin Nutr* **69**, 842-856.
- Vieth R (2006) Critique of the Considerations for Establishing the Tolerable Upper Intake Level for Vitamin D: Critical Need for Revision Upwards. *J Nutr* **136**, 1117-1122.
- Vieth R, Ladak Y & Walfish PG (2003) Age-Related Changes in the 25-Hydroxyvitamin D Versus Parathyroid Hormone Relationship Suggest a Different Reason Why Older Adults Require More Vitamin D. *J Clin Endocrinol Metab* **88**, 185-191.
- Wactawski-Wende J, Kotchen JM, Anderson GL, *et al.* (2006) Calcium plus Vitamin D Supplementation and the Risk of Colorectal Cancer. *N Engl J Med* **354**, 684-696.
- Walters JR, Barley NF, Khanji M & Rhodes-Kendler O (2004) Duodenal expression of the epithelial calcium transporter gene TRPV6: is there evidence for Vitamin D-dependence in humans? *J Steroid Biochem Mol Biol* **89-90**, 317-319.
- Walters JR, Balesaria S, Chavele KM, *et al.* (2006) Calcium Channel TRPV6 Expression in Human Duodenum: Different Relationships to the Vitamin D System and Aging in

Men and Women. *Journal of Bone and Mineral Research* **21**, 1770-1777.

Webb AR & Holick MF (1988) The role of sunlight in the cutaneous production of vitamin D3. *Annu Rev Nutr* **8**, 375-399.

Wortsman J, Matsuoka LY, Chen TC, Lu Z & Holick MF (2000) Decreased bioavailability of vitamin D in obesity. *Am J Clin Nutr* **72**, 690-693.

Zamboni G, Soffiati M, Giavarina D & Tato L (1988) Mineral metabolism in obese children. *Acta Paediatr Scand* **77**, 741-746.

Zehnder D, Bland R, Chana RS, Wheeler DC, Howie AJ, Williams MC, Stewart PM & Hewison M (2002) Synthesis of 1,25-Dihydroxyvitamin D3 by Human Endothelial Cells Is Regulated by Inflammatory Cytokines: A Novel Autocrine Determinant of Vascular Cell Adhesion. *J Am Soc Nephrol* **13**, 621-629.

Zehnder D, Bland R, Williams MC, McNinch RW, Howie AJ, Stewart PM & Hewison M (2001) Extrarenal Expression of 25-Hydroxyvitamin D3-1 α -Hydroxylase. *J Clin Endocrinol Metab* **86**, 888-894.

Ziegler EE, Hollis BW, Nelson SE & Jeter JM (2006) Vitamin D Deficiency in Breastfed Infants in Iowa. *Pediatrics* **118**, 603-610.

Zipitis CS, Markides GA & Swann IL (2006) Vitamin D deficiency: prevention or treatment? *Arch Dis Child* **91**, 1011-1014

Zittermann A, Schleithoff SS & Koerfer R (2005) Putting cardiovascular disease and vitamin D insufficiency into perspective. *Br J Nutr* **94**, 483-492.